Europäisches Patentamt

European Patent Office

Office européen des brevets



11 Publication number:

0 510 773 A1

(P)

EUROPEAN PATENT APPLICATION

2) Application number: 92201136.6

② Date of filing: 22.04.92

(5) Int. Cl.⁵: **C12N 15/50**, A61K 39/215, G01N 33/569, C07K 15/00

- Priority: 25.04.91 EP 91303737
- Date of publication of application:
 28.10.92 Bulletin 92/44
- Designated Contracting States:
 AT BE CH DE DK ES FR GB GR IT LI LU MC
 NL PT SE
- 7) Applicant: AKZO N.V. Velperweg 76 NL-6824 BM Amhem(NL)
- 2 Inventor: Brown, Thomas David Kay
 The Ferns, 8 High Street, Needingworth
 Huntingdon, Cambs. PE 1735A(GB)
 Inventor: Horsburgh, Brian Colin
 c/o Darwin College, Silver Street
 Cambridge CB3 9EV(GB)
- Representative: Hermans, Franciscus G.M. et al P.O. Box 20 NL-5340 BH Oss(NL)
- (a) Canine coronavirus subunit vaccine.
- The invention is related to a nucleic acid sequence encoding a Canine coronavirus (CCV) spike protein. Such a protein can be used for the immunization of dogs against CCV infection. The nucleic acid sequence encoding the CCV spike protein can be applied for the preparation of the spike protein by means of genetic engineering techniques or can be applied for the preparation of vector vaccines.

The present invention is concerned with a nucleic acid sequence encoding a CCV spike protein, a recombinant vector or recombinant vector virus comprising such a nucleic acid sequence, a host cell transformed with such a recombinant vector or infected with the recombinant vector virus, as well as a vaccine against CCV infection in dogs.

Canine coronavirus (CCV) is a member of the distinct viral family of Coronavirus. Viruses belonging to this genus are known to infect a variety of animal species including man. They cause diverse diseases, such as gastro-enteritis (in swine, turkeys, mice, calves, dogs, cats and man), sativary gland infection (in rodents), respiratory disease (in man, swine, avians and dogs) and encephalitis (in young swine).

CCV was first isolated from military dogs in Germany in 1971 and has found to be highly contagious and it spreads rapidly among susceptible dogs. Usually, the CCV is ingested on materials contaminated by infectious feces. Oral infection leads to viral replication in epithelial cells of the small intestine and CCV has also been found in the intestinal lymph nodes.

The signs of the disease can develop 1-3 days following infection and include vomiting, diarrhoea, anorexia, depression and dehydration. The persistence and severity of signs is often related to stress and the presence of other viruses, parasites or bacteria. Whereas the enteric symptoms are dominant, respiratory signs including nasal and ocular discharge have also been reported.

Dogs are the only known host of the CCV. Although CCV inoculation of cats and pigs results in infection, no clinical disease will be caused by CCV in these species. There is no evidence that humans, cattle and mice are susceptible to CCV.

Cross protection studies have shown that the Coronaviruses induce little or no immunity to each other. For example, experimental infection of dogs with transmissible gastro-enteritis virus (TGEV) of pigs or feline infectious peritonitis virus (FIPV) of cat does not protect them against the effects of a subsequent CCV infection.

Coronaviruses consist of a group of enveloped viruses containing a genome consisting of a single-stranded RNA of about 30 kb. This genome encodes inter alia three important structural proteins: a spike protein (S), a membrane protein (M) and a nucleocapsid protein (N). The glycosylated spike protein S_0 is cleaved to form S_1 and S_2 in some coronaviruses. Two or three copies of each of S_1 and S_2 form a characteristic CCV surface structure, the spike or peplomer. The spike protein and its constituent polypeptides thereof play an important role in inducing a virus neutralizing immune response in infected animals.

Conventional CCV vaccines comprise chemically inactivated virus vaccines or modified live-virus vaccines. However, inactivated vaccines require additional immunizations, disadvantageously contain adjuvants and are expensive to produce. Further, some infectious virus particles may survive the inactivation process and may cause disease after administration to the animal.

In general, attenuated live virus vaccines are preferred because they evoke an immune response often based on both humoral and cellular reactions. Up to now, such vaccines based on CCV strains can only be prepared by serial passage of virulent strains in tissue culture. However, because of this treatment uncontrolled mutations are introduced into the viral genome, resulting in a population of virus particles heterogeneous in their virulence and immunizing properties. In addition it is well known that such traditional attenuated live virus vaccines can revert to virulence resulting in disease of the inoculated animals and the possible spread of the pathogen to other animals.

Improved vaccines might be constructed, based on recombinant DNA technology, which only contain the necessary and relevant CCV immunogenic material capable of eliciting an immune response against the CCV pathogens, or which contain the genetic information encoding said material, and do not display abovementioned disadvantages of the live or inactivated vaccines.

According to the present invention, an isolated and purified nucleic acid sequence encoding a polypeptide having one or more immunogenic determinants of a CCV spike protein is provided which can be applied for the preparation of a vaccine for the immunization of dogs against CCV infection.

"Nucleic acid sequence" as used herein refers to a polymeric form of nucleotides of any length, both to ribonucleic acid sequences and to deoxy ribonucleic acid sequences. In principle, this term refers to the primary structure of the molecule. Thus, this term includes double and single stranded DNA, as well as double and single stranded RNA, and modifications thereof.

In general, the term "polypeptide" refers to a molecular chain of amino acids with a biological activity, does not refer to a specific length of the product and if required can be modified in vivo or in vitro, for example by glycosylation, amidation, carboxylation or phosphorylation; thus inter alia, peptides, oligopeptides and proteins are included.

The term "polypeptide having one or more immunogenic determinants of a CCV spike protein" refers to a polypeptide having one or more epitopes capable of eliciting a protective immune response in a dog against CCV infection or disease.

5

30

45

In particular, the present invention provides a nucleic acid sequence encoding a polypeptide having one or more immunogenic determinants of the CCV spike protein which has an amino acid sequence shown in SEQ ID NO: 2, 4 or 6.

Also included within the scope of the present invention are nucleic acid sequences encoding a functional variant of the polypeptide shown in SEQ ID NO: 2, 4 or 6. These functional variants are polypeptides having an amino acid sequence derived from the amino acid sequence specifically disclosed in SEQ ID NO: 2, 4 or 6 but retain one or more immunogenic determinants of a CCV spike protein, i.e. said variants having one or more epitopes capable of eliciting a protective immune response in a dog against CCV infection or disease.

It will be understood that for the particular polypeptide embraced herein, derived from the CCV-6, Insavc-1 or Liverpool C54 strain, natural variations can exist between individual viruses or strains of canine coronaviruses. These variations may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of (an) amino acid(s) in said sequence. Amino acid substitutions from which can be expected that they do not essentially alter biological and immunological activities, have been described. Amino acid replacements between related amino acids or replacements which have occurred frequently in evolution are, inter alia Ser/Ala, Ser/Gly, Asp/Gly, Asp/Asn, Ile/Val (see Dayhof, M.D., Atlas of protein sequence and structure, Nat. Biomed. Res. Found., Washington D.C., 1978, vol. 5, suppl. 3). Based on this information Lipman and Pearson developed a method for rapid and sensitive protein comparison (Science 227, 1435-1441, 1985) and determining the functional similarity between homologous polypeptides. Nucleic acid sequences encoding such homologous functional variants are included within the scope of this invention. Moreover, the potential exists to use recombinant DNA technology for the preparation of nucleic acid sequences encoding these various functional variants.

Nucleic acid sequences according to the invention may be derived from isolates of CCV strains such as CCV-6, Insavc-1 (EP 396,193), CCV 1-71 (ATCC VR-809) or CCV TN449 (ATCC VR-2068).

In another aspect of the invention nucleic acid sequences described above are provided which can beused for the preparation of a vaccine to protect cats against FIPV infection.

The information provided in SEQ ID NO: 1-6 allows a person skilled in the art to isolate and identify the nucleic acid sequences encoding the various functional variant polypeptidec mentioned-above having corresponding immunological characteristics with the CCV spike protein specifically disclosed herein. The generally applied Southern blotting technique or colony hybridization can be used for that purpose' (Experiments in Molecular Biology, ed. R.J. Slater, Clifton, U.S.A., 1986; Singer-Sam, J. et al., Proc. Natl. Acad. Sci. 80, 802-806, 1983; Maniatis T. et al., Molecular Cloning, A laboratory Manual, second edition, Cold Spring Harbor Laboratory Press, USA, 1989). For example, RNA or cDNA derived from a specific CCV strain is electrophoresed and transferred, or "blotted" thereafter onto a piece of nitrocellulose filter. It is now possible to identify CCV spike protein nucleic acid sequences on the filter by hybridization to a defined labeled DNA fragment or "probe", i.e. a (synthetic) poly- or oligonucleotide sequence fragment of the nucleic acid sequence shown in SEQ ID NO: 1, 3 or 5 which under specific conditions of salt concentration and temperature hybridizes to the homologous nucleic acid sequences present on the filter. After washing the filter, hybridized material may be detected by autoradiography. The corresponding DNA fragment can now be eluted from the agarose gel and used to direct the synthesis of a functional variant of the polypeptide disclosed in SEQ ID NO: 2, 4 or 6.

Therefore, a preferred functional variant according to the invention is a polypeptide comprising one or more immunogenic determinants of a CCV spike protein and is encoded by a nucleic acid sequence which hybridizes to the DNA sequence shown in SEQ ID NO: 1, 3 or 5.

In another way CCV cDNA may be cloned into a kgt11 phage as described by Huynh et al. (In: D. Glover (ed.), DNA Cloning: A Practical Approach, IRL Press Oxford, 49-78, 1985) and expressed in a bacterial host. Recombinant phages can then be screened with polyclonal serum raised against the purified CCV spike protein disclosed in SEQ ID NO: 2, 4 or 6 determining the presence of corresponding immunological regions of the variant polypeptide. The production of the polyclonal serum to be used herein elicited against the CCV spike protein is described below.

As is well known in the art, the degeneracy of the genetic code permits substitution of bases in a codon resulting in an other codon but still coding for the same amino acid, e.g. the codon for the amino acid glutamic acid is both GAT and GAA. Consequently, it is clear that for the expression of a polypeptide with the amino acid sequence shown in SEQ ID NO: 2, 4 or 6 use can be made of a derivate nucleic acid sequence with such an alternative codon composition different from the nucleic acid sequence shown in SEQ ID NO: 1, 3 or 5, respectively.

Furthermore, also fragments of the nucleic acid sequences encoding the specifically disclosed CCV

spike protein or functional variants thereof as mentioned above are included in the present invention.

The term "fragment" as used herein means a DNA or amino acid sequence comprising a subsequence of the nucleic acid sequence or polypeptide of the invention. Said fragment is or encodes a polypeptide having one or more immunogenic determinants of a CCV spike protein, i.e. has one or more epitopes which are capable of eliciting a protective immune response in a dog. Methods for determining usable polypeptide fragments are outlined below. Fragments can inter alia be produced by enzymatic cleavage of precursor molecules, using restriction endonucleases for the DNA and proteases for the polypeptides. Other methods include chemical synthesis of the fragments or the expression of polypeptide fragments by DNA fragments.

Typical sequences encoding the CCV spike protein precursor are shown in SEQ ID NO: 1, 3 and 5. These cDNA sequences are about 4328, 4352 and 4358 nucleotides in length, respectively, and encode a polypeptide of 1443, 1451 and 1453 amino acids, respectively.

A preferred nucleic acid sequence according to the invention is characterized in that said sequence contains at least part of the DNA sequence disclosed in SEQ ID NO: 1, 3 or 5.

A nucleic acid sequence according to the invention may be isolated from a particular CCV strain and multiplied by recombinant DNA techniques including polymerase chain reaction (PCR) technology or may be chemically synthesized in vitro by techniques known in the art.

All modifications resulting in the above-mentioned functional variants of the specifically exemplified polypeptide are included within the scope of the present invention for as long as the resulting polypeptides retain one or more immunogenic determinants of a CCV spike protein.

A nucleic acid sequence according to the present invention can be ligated to various replication effecting DNA sequences with which it is not associated or linked in nature resulting in a so called recombinant vector molecule which can be used for the transformation of a suitable host. Useful recombinant vector molecules, are preferably derived from, for example plasmids, bacteriophages, cosmids or viruses.

Specific vectors or cloning vehicles which can be used to clone nucleic acid sequences according to the invention are known in the art and include inter alia plasmid vectors such as pBR322, the various pUC, pGEM and Bluescript plasmids, bacteriophages, e.g. kgt-Wes,Charon 28 and the M13 derived phages or viral vectors such as SV40, adenovirus or polyoma virus (see also Rodriquez, R.L. and D.T. Denhardt, ed., Vectors: A survey of molecular cloning vectors and their uses, Butterworths, 1988; Lenstra, J.A. et al., Arch. Virol. 110, 1-24, 1990). The methods to be used for the construction of a recombinant vector molecule according to the invention are known to those of ordinary skill in the art and are inter alia set forth in Maniatis, T. et al. (Molecular Cloning A Laboratory Manual, second edition; Cold Spring Harbor Laboratory, 1989).

For example, the insertion of the nucleic acid sequence according to the invention into a cloning vector can easily be achieved when both the genes and the desired cloning vehicle have been cut with the same restriction enzyme(s) as complementary DNA termini are thereby produced.

Alternatively, it may be necessary to modify the restriction sites that are produced into blunt ends either by digesting the single-stranded DNA or by filling in the single-stranded termini with an appropriate DNA polymerase. Subsequently, blunt end ligation with an enzyme such as T4 DNA ligase may be carried out.

If desired, any restriction site may be produced by ligating linkers onto the DNA termini. Such linkers may comprise specific oligonucleotide sequences that encode restriction site sequences. The restriction enzyme cleaved vector and nucleic acid sequence may also be modified by homopolymeric tailing.

"Transformation", as used herein, refers to the introduction of a heterologous nucleic acid sequence into a host cell, irrespective of the method used, for example direct uptake or transduction. The heterologous nucleic acid sequence may be maintained through autonomous replication or alternatively, may be integrated into the host genome. If desired, the recombinant vector molecules are provided with appropriate control sequences compatible with the designated host which can regulate the expression of the inserted nucleic acid sequence. In addition to microorganisms, culture of cells derived from multicellular organisms may also be used as hosts.

The recombinant vector molecules according to the invention preferably contain one or more marker activities that may be used to select for desired transformants, such as ampicillin and tetracycline resistance in pBR322, ampicillin resistance and β -galactosidase activity in pUC8.

A suitable host cell is a microorganism or cell which can be transformed by a nucleic acid sequence encoding a polypeptide or by a recombinant vector molecule comprising such a nucleic acid sequence and which can if desired be used to express said polypeptide encoded by said nucleic acid sequence. The host cell can be of procaryotic origin, e.g. bacteria such as Escherichia coli, Bacillus subtilis and Pseudomonas species; or of eucaryotic origin such as yeasts, e.g. Saccharomyces cerevisiae or higher eucaryotic cells such as insect, plant or mammalian cells, including HeLa cells and Chinese hamster ovary (CHO) cells.

50

20

Insect cells include the Sf9 cell line of Spodoptera frugiperda (Luckow et al., Bio-technology 6, 47-55, 1988). Information with respect to the cloning and expression of the nucleic acid sequence of the present invention in eucaryotic cloning systems can be found in Esser, K. et al. (Plasmids of Eukaryotes, Springer-Verlag, 1986).

In general, prokaryotes are preferred for the construction of the recombinant vector molecules useful in the invention. For example E.coli K12 strains are particularly useful such as DH5 α or JM101.

For expression nucleic acid sequences of the present invention are introduced into an expression vector, i.e. said sequences are operably linked to expression control sequences. Such control sequences may comprise promoters, enhancers, operators, inducers, ribosome binding sites etc. Therefore, the present invention provides a recombinant vector molecule comprising a nucleic acid sequence encoding the CCV spike protein operably linked to expression control sequences, capable of expressing the DNA sequences contained therein in (a) transformed host cell(s).

It should, of course, be understood that the nucleotide sequences inserted at the selected site of the cloning vector may include nucleotides which are not part of the actual structural gene for the desired polypeptide or may include only a fragment of the complete structural gene for the desired protein as long as transformed host will produce a polypeptide having at least one or more immunogenic determinants of a CCV spike protein.

When the host cells are bacteria, illustrative useful expression control sequences include the Trp promoter and operator (Goeddel, et al., Nucl. Acids Res. 8, 4057, 1980); the lac promoter and operator (Chang, et al., Nature 275, 615, 1978); the outer membrane protein promoter (Nakamura, K. and Inouge, Mc, EMBO J. 1, 771-775, 1982); the bacteriophage k promoters and operators (Remaut, E. et al., Nucl. Acids Res. 11, 4677-4688, 1983); the α-amylase (B. subtilis) promoter and operator, termination sequence and other expression enhancement and control sequences compatible with the selected host cell. When there we host cell is yeast, illustrative useful expression control sequences include, e.g., α-mating factor. For insect cells the polyhedrin or p10 promoters of baculoviruses can be used (Smith, G.E. et al., Mol. Cell. Biol. 3, 2156-65, 1983). When the host cell is of mammalian origin illustrative useful expression control sequences include, e.g., the SV-40 promoter (Berman, P.W. et al., Science 222, 524-527, 1983) or, e.g. the metallothionein promoter (Brinster, R.L., Nature 296, 39-42, 1982) or a heat shock promoter (Voellmy et al., Proc. Natl. Acad. Sci. USA 82, 4949-53, 1985). For maximizing gene expression, see also Roberts and Lauer (Methods in Enzymology 68, 473, 1979).

Therefore, the invention also comprises (a) host cell(s) transformed with a nucleic acid sequence or recombinant expression vector molecule described above, capable of producing the CCV spike protein by expression of the nucleic acid sequence.

The present invention also provides a process for the preparation of a purified polypeptide displaying immunological characteristics of a CCV spike protein, i.e. the polypeptide has one or more immunogenic determinants of a CCV spike protein, essentially free from whole viruses or other protein with which it is ordinarily associated.

More particularly, the invention provides a process for the preparation of a polypeptide comprising at least part of the amino acid sequence shown in SEQ ID NO: 2, 4 or 6 or a functional variant thereof.

In addition a polypeptide substantially comprising an immunogenic fragment of the CCV spike protein which can be used for immunization of dogs against CCV infection or diagnostic purposes, is prepared in the present invention. Various methods are known for detecting such usable immunogenic fragments within an amino acid sequence.

Suitable immunochemically active polypeptide fragments of a polypeptide according to the invention containing (an) epitope(s) can be found by means of the method described in Patent Application WO 86/06487, Geysen, H.M. et al. (Prod. Natl. Acad. Sci. 81, 3998-4002, 1984), Geysen, H.M. et al. (J. Immunol. Meth. 102, 259-274, 1987) based on the so-called pep-scan method, wherein a series of partially overlapping peptides corresponding with partial sequences of the complete polypeptide under consideration, are synthesized and their reactivity with antibodies is investigated.

In addition, a number of regions of the polypeptide, with the stated amino acid sequence, can be designated epitopes on the basis of theoretical considerations and structural agreement with epitopes which are now known. The determination of these regions is based on a combination of the hydrophilicity criteria according to Hopp and Woods (Proc. Natl. Acad. Sci. 78, 3824-3828, 1981) and the secondary structure aspects according to Chou and Fasman (Advances in Enzymology 47, 45-148, 1987).

T-cell epitopes which may be necessary can likewise be derived on theoretical grounds, e.g. with the aid of Berzofsky's amphiphilicity criterion (Science 235, 1059-62, 1987).

In another embodiment of the invention a polypeptide having an amino acid sequence encoded by a nucleic acid sequence mentioned above is used.

Immunization of dogs against CCV infection can, for example be achieved by administering to the animals a polypeptide prepared according to the process mentioned above in an immunologically relevant context as a so-called subunit vaccine. The subunit vaccine according to the invention may comprise a polypeptide in a pure form, optionally in the presence of a pharmaceutically acceptable carrier. The polypeptide can optionally be covalently bound to a non-related protein, which, for example can be of advantage in the purification of the fusion product. Examples are β -galactosidase, protein A, prochymosine, blood clotting factor Xa, etc.

In some cases the ability to raise neutralizing antibodies against these polypeptides per se may be low. Small fragments are preferably conjugated to carrier molecules in order to raise their immunogenicity. Suitable carriers for this purpose are macromolecules, such as natural polymers (proteins like key hole limpet hemocyanin, albumin, toxins), synthetic polymers like polyamino acids (polylysine, polyalanine), or micelles of amphiphilic compounds like saponins. Alternatively these fragments may be provided as polymers thereof, preferably linear polymers.

Polypeptides to be used in such subunit vaccines can be prepared by methods known in the art, e.g. by isolating said polypeptides from CCV, by recombinant DNA techniques or by chemical synthesis.

If required these polypeptides to be used in a vaccine can be modified in vitro or in vivo, for example by glycosylation, amidation, carboxylation or phosphorylation.

An alternative to subunit vaccines are five vector vaccines. A nucleic acid sequence according to the invention is introduced by recombinant DNA techniques into a micro-organism (e.g. a bacterium or virus) in such a way that the recombinant micro-organism is still able to replicate thereby expressing a polypeptide coded by the inserted nucleic acid sequence and eliciting an immune response in the infected host animal.

A preferred embodiment of the present invention is a recombinant vector virus comprising a heterologous nucleic acid sequence described above, capable of expressing the DNA sequence in (a) host cell(s) or host animal infected with the recombinant vector virus. The term "heterologous" indicates that the nucleic acid sequence according to the invention is not normally present in nature in the vector virus.

Furthermore, the invention also comprises (a) host cell(s) or cell culture infected with the recombinant vector virus, capable of producing the CCV protein by expression of the nucleic acid sequence.

For example the well known technique of in vivo homologous recombination can be used to introduce a heterologous nucleic acid sequence, e.g. a nucleic acid sequence according to the invention into the genome of the vector virus.

First, a DNA fragment corresponding with an insertion region of the vector genome, i.e. a region which can be used for the incorporation of a heterologous sequence without disrupting essential functions of the vector such as those necessary for infection or replication, is inserted into a cloning vector according to standard recDNA techniques. Insertion-regions have been reported for a large number of micro-organisms (e.g. EP 80,806, EP 110,385, EP 83,286, US 4,769,330 and US 4,722,848).

Second, if desired, a deletion can be introduced into the insertion region present in the recombinant vector molecule obtained from the first step. This can be achieved for example by appropriate exonuclease III digestion or restriction enzyme treatment of the recombinant vector molecule from the first step.

Third, the heterologous nucleic acid sequence is inserted into the insertion-region present in the recombinant vector molecule of the first step or in place of the DNA deleted from said recombinant vector molecule. The insertion region DNA sequence should be of appropriate length as to allow homologous recombination with the vector genome to occur. Thereafter, suitable cells can be infected with wild-type vector virus or transformed with vector genomic DNA in the presence of the recombinant vector molecule containing the insertion flanked by appropriate vector DNA sequences whereby recombination occurs between the corresponding regions in the recombinant vector molecule and the vector genome. Recombinant vector progeny can now be produced in cell culture and can be selected for example genotypically or phenotypically, e.g. by hybridization, detecting enzyme activity encoded by a gene co-integrated along with the heterologous nucleic acid sequence, or detecting the antigenic heterologous polypeptide expressed by the recombinant vector immunologically.

Next, this recombinant micro-organism can be administered to the dogs for immunization whereafter it maintains itself for some time, or even replicates in the body of the inoculated animal, expressing in vivo a polypeptide coded for by the inserted nucleic acid sequence according to the invention resulting in the stimulation of the immune system of the inoculated animal. Suitable vectors for the incorporation of a nucleic acid sequence according to the invention can be derived from viruses such as pox viruses, e.g. vaccinia virus (EP 110,385, EP 83,286, US 4,769,330 and US 4,722;848), herpes viruses such as Feline Herpes virus, (canine) adeno virus (WO 91/11525) or inflüenza virus, or bacteria such as E. coli or specific Salmonella species. With recombinant microorganisms of this type, the polypeptide synthesized in the host can be exposed as a cell surface antigen. In this context fusion of the said polypeptide with OMP proteins,

or pilus proteins of for example E. coli or synthetic provision of signal and anchor sequences which are recognized by the organism are conceivable. It is also possible that the sald immunogenic polypeptide, if desired as part of a larger whole, is released inside the animal to be immunized. In all of these cases it is also possible that one or more immunogenic products will find expression which generate protection against various pathogens and/or against various antigens of a given pathogen.

A vaccine according to the invention can be prepared by culturing a host cell infected with a recombinant vector virus comprising a nucleic acid sequence according to the invention, whereafter virus containing cells and/or recombinant vector viruses grown in the cells can be collected, optionally in a pure form, and formed to a vaccine optionally in a lyophilized form.

Host cells transformed with a recombinant vector molecule according to the invention can also be cultured under conditions which are favourable for the expression of a polypeptide coded by said nucleic acid sequence. Vaccines may be prepared using samples of the crude culture, host cell lysates or host cell extracts, although in another embodiment more purified polypeptides according to the invention are formed to a vaccine, depending on its intended use. In order to purify the polypeptides produced, host cells transformed with a recombinant vector according to the invention are cultured in an adequate volume and the polypeptides produced are isolated from such cells or from the medium if the protein is excreted. Polypeptides excreted into the medium can be isolated and purified by standard techniques, e.g. salt fractionation, centrifugation, ultrafiltration, chromatography, gel filtration or immuno affinity chromatography, whereas intra cellular polypeptides can be isolated by first collecting said cells, disrupting the cells, for example by sonication or by other mechanically disruptive means such as French press followed by separation of the polypeptides from the other intra cellular components and forming the polypeptides to a vaccine. Cell disruption could also be accomplished by chemical (e.g. EDTA treatment) or enzymatic means, such as lysozyme digestion.

The vaccine according to the invention can be administered in a convential active immunization scheme: single or repeated administration in a manner compatible with the dosage formulation and in such amount as will be prophylactically and/or therapeutically effective and immunogenic, i.e. the amount of immunizing antigen or recombinant micro-organism capable of expressing said antigen that will induce immunity in a dog against challenge by a virulent CCV. Immunity is defined as the induction of a significant level of protection in a population of dogs after vaccination compared to an unvaccinated group.

For live viral vector vaccines the dose rate per dog may range from 10^5 - 10^8 pfu.

A typical subunit vaccine according to the invention comprises 10 μg - 1 mg of the polypeptide according to the invention.

The administration of the vaccine can be done, e.g. intradermally, subcutaneously, intramuscularly, intraperitonially, intravenously, orally or intranasally.

Additionally the vaccine may also contain an aqueous medium or a water containing suspension, often mixed with other constituents, e.g. in order to increase the activity and/or shelf life. These constituents may be salts, pH buffers, stabilizers (such as skimmed milk or casein hydrolysate), emulsifiers adjuvants to improve the immune response (e.g. oils, muramyl dipeptide, aluminiumhydroxide, saponin, polyanions and amphipatic substances) and preservatives.

It is clear that a vaccine according to the invention may also contain immunogens related to other pathogens of dogs or may contain nucleic acid sequences encoding these immunogens, like antigens of Canine parvovirus (CPV), Canine Distemper virus, Canine Adenovirus I, Canine Adenovirus II, Canine Parainfluenza virus, Canine Rotavirus or Leptospira canicola to produce a multivalent vaccine.

Example 1

10

30

35

50

55

A.

1. Preparation of genomic viral RNA of CCV-6 and Liverpool C54 strain

Confluent A-72 cells grown in plastic tissue culture flasks using the Wellcome modification of minimal Eagle's medium (MEM) and 10% foetal bovine serum were infected with CCV (NVSL Challenge virus CCV-6 from the National Veterinary Service Laboratory, PO Box 844, Ames, Iowa 50010, USA) at a multiplicity of infection (MOI) of approximately 0.1. After 24 h the culture supernatant was harvested, chilled to 4 °C and cell debris removed by centrifugation at 3000 x g for 15 min. Virus was pelletted from the supernatant at 53.000 x g for 2 h in a Beckman type 19 rotor. The pellet was resuspended in 5 ml of TNE (10 mM Tris-Cl, 100 mM NaCl, 1 mM EDTA, pH 7.5) using a Dounce homogeniser and layered onto a 32 ml linear 20-60% gradient of sucrose in TNE. The virus was banded isopycnically by overnight centrifugation at 100.000 x g in a Beckman SW28 rotor. The gradient was fractionated and the A280's and densities of the fractions determined. A peak was identified at the characteristic

density of 1.18 g/cc. The peak fractions were pooled, diluted in TNE and the putative virus pelletted by centrifugation at $100.000 \times g$ for 2 h in the Beckman SW28 rotor. RNA was isolated from the virus pellet using two approaches:

A. The pellet was resuspended in 0.1 M Tris-Cl pH 8.0 containing 0.1% SDS and digested for 3 h at 50 °C with 20 µg/ml of proteinase K. The mixture was deproteinised using phenol:chloroform:isoamyl alcohol (50:49:1) saturated with TE (10 mM Tris-Cl 1 mM EDTA) and the nucleic acid recovered by precipitation with 2.5 volumes of ethanol/0.3 M sodium acetate pH 5.2. The preparation was analysed on a Tris-borate EDTA 1% agarose gel containing 0.1% SDS; a high molecular weight RNA band was identified with the characteristic mobility of coronavirus genomic RNA.

B. The virus pellet was homogenised in 6 M guanidinium isothiocyanate/5 mM sodium citrate (pH 7.0)/0.1 M mercaptoethanol/0.5% N-lauroyl sarcosinate and 1 g/ml of CsCl added to each 2.5 ml of the homogenate. The mixture was then layered onto a 5.7 M CsCl/0.1 M EDTA pad and centrifuged at 108.000 x g for 12 h at 20 °C. The pellet of RNA was dissolved in TE containing 0.1% SDS. The preparation was analysed as described above.

2. cDNA cloning of CCV genomic RNA

First strand synthesis from 2 µg of CCV genomic RNA prepared as described in 1A above was primed with 1 ng of a specific oligonucleotide (5' TTTTCAAATTGTCTTCTACTT 3') using 40 units of AMV reverse transcriptase in a reaction volume of 25 μl containing 20 mM Tris-Cl (pH 8.3 at 42 °C), 0.14 M KCl, 10 mM MgCl₂, 1 mM dNTP's, 4 mM dithiothreitol, 25 units of human placental ribonuclease inhibitor. The reaction mixture was incubated for 1 h at 42 °C. Second strand synthesis was achieved by addition of 46 µl of a reaction mixture containing 7.6 mM MgCl2, 0.109 M Tris-Cl pH 7.4, 16.3 mM (NH₄)₂SO₄, 1000 units/ml RNaseH, 10.000 units/ml E, coli DNA polymerase 1 to the first strand reaction and incubation at 12 °C for 1 h followed by incubation at 22 °C for a further 1 h. The reaction products were deproteinised by two extractions with phenol:chloroform:isoamyl alcohol (50:49:1) saturated with TE and precipitated with 2 volumes of ethanol/0.3 M sodium acetate pH 5.2. The cDNA was tailed with C residues using terminal deoxynucleotidyl transferase using the buffer and conditions supplied by the manufacturer (Bethesda Research Laboratories, Gaithersburg, Maryland 20877, USA). It was then size fractionated on a 2 ml Sephacryl S-1000 column and cDNA of size greater than 500 base pairs pooled, ethanol precipitated and dissolved in TE. 50 ng of this cDNA was annealed with 250 ng of dG-tailed Pstl cut pUC119. The mixture was transformed into E. coli TG-1. Ampicillin resistant transformants were picked and screened for CCV cDNA inserts using a cDNA probe produced by random priming of reverse transcription from CCV genomic RNA. Positive colonies were screened for size of cDNA inserts by Pstl digestion of mini-prep DNA. The relationships between inserts were established by restriction enzyme analysis. The clone pBHI was selected for sequence analysis. The size of the pBHI insert (4.0 kb) was insufficient to cover the complete CCV spike coding region and a further round of cDNA synthesis and cloning was carried out using another specific primer (5' CTAGGTAGTAACAC 3'). The RNA used was isolated as described in 1B above. cDNA synthesis was achieved using a Boehringer Mannheim (Boehringer Mannheim UK, Bell Lane, Lewes, East Sussex BN7 1LG) cDNA synthesis kit according to the manufacturer's instructions. In summary first strand synthesis was again achieved using AMV reverse transcriptase, second strand synthesis by the action of E. coli DNA polymerase 1 and RNaseH. The cDNA was blunt ended by the action of T4 DNA polymerase. The cDNA was ligated into Smal-cut phosphatased pUC18 using T4 DNA ligase and the DNA transformed into E. coli TG1. Ampicillin resistant clones were initially screened for inserts using blue/white selection on X-gal (5-bromo-4-chloro-3-indolyl-8-Dgalactopyranoside) plates. White colonies were picked and screened for the presence of CCV cDNA inserts as described above. Clone pBH2 (size 2.8 kb) was selected for sequence analysis. The same strategy as outlined in Example 1.1.B. and 1.2. for CCV strain CCV6 was carried out for the isolation of the spike gene of the CCV C54 strain. Three overlapping clones, pBH3, pBH4 and pBH11 covered the spike gene to the blunt end.

3. DNA sequencing The cDNA inserts from clones pBH1, pBH2, and pBH3, pBH4 and pBH11 were sequenced using the Sanger dideoxy chain termination method. This shotgun approach was supplemented as necessary with sequencing from specific oligonucleotide primers on double stranded plasmid DNA templates. For the shotgun analysis insert DNA was excised from the vector sequences, circularised, sonicated, size selected on agarose gels and cloned into Smal-cut, phosphatased M13mp8. Shotgun sequence data were assembled using the DBUTIL and DBAUTO programs of Staden and analysed using the ANALYSEQ/NIP packages of Staden. A VAX 8350 and micro VAX 3100 (Digital Equipment Corporation) were used. The sequence data are presented in SEQ ID NO: 1

5

10

15

20

25

30

35

40

45

50

and 5.

В.

5

10

15

20

25

30

35

40

45

55

1. Preparation of genomic viral RNA of Insavc-1 strain

Confluent A-72 cells grown in plastic tissue culture flasks using the Wellcome modification of minimal Eagle's medium (MEM) and 10% F.C.S. were infected with CCV strain Insavc-1 (Bert) (Intervet Labs.) at a m.o.i. of approximately 0.1. After 48 hours the culture supernatant was harvested, chilled to 4 ° and cell debris removed by centrifugation at 3000xg for 15 minutes. Virus was pelleted from the supernatant at 53000xg for 2 hrs in a Beckman type 19 rotor. The virus pellet was homogenized in 3.5 mls of 6M guanidinium isothiocyanate/5mM sodium citrate (pH 7.0), 0.1M mercaptoethanol, 0.5% N-lauroyl sarcosinate.

The homogenate was layered onto a 5.7 M CsCl pad (1 ml) and centrifuged at 108000 g for 18 hours at 18 °C. The pellet of RNA was dissolved in TE containing 0.1% SDS, then precipitated twice with 2.5 volumes of ethanol/0.3M NaOAc pH 5.2. The preparation was analysed as a Tris-borate EDTA 1% agarose gel containing 0.1% SDS, high molecular weight RNA band was identified with the characteristic mobility of coronavirus genomic RNA.

2. cDNA and PCR cloning of CCV genomic RNA

First and second strand synthesis from 2 µg of CCV genomic RNA prepared as aforementioned was primed with oligo dT and random pentanucleotides from the Boehringer cDNA synthesis kit under the conditions specified by the manufacturers protocol.

The resultant blunt ended cDNA produced from this reaction was ligated into Sma1-cut-phosphatased pUC 119 using T4 DNA ligase and the DNA transformed into E. coli TG-1. Ampicillin resistant clones were initially screened for inserts using blue/white selection on x-gal (5-bromo-4-chloro-3-indetyl-B-D-galaclopyramoside) plates. White colonies were picked and screened for the presence of CCV cDNA inserts using randomly primed CCV RNA as a probe. Five positive clones were identified.

Plasmid pBH6 was generated using the polymerase chain reaction (PCR). Sequence information from the ends of pBH5 and pBH7 allowed the design of primers BH7 and BH8. A Not 1 site was incorporated into the oligo's to facilitate cloning. Briefly, approximately 1 ng of first-strand reaction as described previously was deproteinized by two extractions with phenol:chloroform:Isoramyl alcohol (50:49:1) saturated with TE, passed down a G50 spin column and precipitated with two volumes of ethanol/0.3 M sodium acetate pH 5.2. The DNA:RNA hybrids were resuspended in 15 µl TE. The PCR reaction was carried out with the Techne programmable Dri-block PHC-1.

The generated fragment was phenol/chloroform ethanol precipitated as before and resuspended.in. 20 µl of TE. The DNA was cleaved with Not 1 under conditions recommended by the enzyme manufacturer, and gel eluted. The Not 1 fragment was then ligated to Not 1 cut phosphatased vector using T4 DNA ligase and the DNA transformed into E. coli TG-1. Clones containing inserts were identified as previously described.

3. DNA sequencing

The cDNA inserts from clones pBH5, pBH7, pBH8, pBH9, pBH10 and the PCR insert pBH6, were sequenced using the Sanger dideoxy chain terminations method as described by Barrell and Bankier (Methods in Enzymology 155, 51-93, 1987). This shotgun approach was supplemented as necessary with sequencing from specific oligonucleotide primers on double stranded or single stranded (f1 origin in pUC 119) plasmid DNA.

For shotgun analyses, insert DNA was excised from the vector sequences, selfligated, sonicated, end-repaired, size selected on 1% agarose gels, cloned into Sma 1-cut phosphatased M13mp18. Shotgun sequence data were assembled and analysed using the SAP programmes of Standen. A Vax 8350 and MicroVax 3100 (Digital Equipment Corporations) were used. The sequence data are presented SEQ ID NO.: 3.

50 Example 2

2.1. Generation of vaccinia virus Vac4b-C6

2.1.1. Assembly of CCV6 full length spike protein gene.

The full length coding region of the S gene of CCV6 was reconstructed from 2 overlapping cDNA clones, BH1 and BH2. The cloning strategy is illustrated in figure 1. The 3.0 kb insert from pBH1 has identity to S and 1b. In order to express S, the polymerase coding sequence had to be removed. The

sequence immediately 5' of the initiating methionine, CTAAACTTTGGTAATCACTTGG TTAATGTGCC ATG was modified by site directed mutagenesis. Four bases, ATCC were looped in between the TGG and TTA bases to create a unique BamHI site, GGATCC. Mutants were screened by restriction enzyme digestion. Positive clones were sequenced across this site as the Klenow fragment of E. coli DNA polymerase used in the mutagenesis reaction can introduce unspecified mutations at a very low frequency. A mutant which had the introduced BamHI site was selected and designated pBHI-bam. This plasmid overlapped pBH2 by approximately 300 bp. A unique AfIII site was located in this region of overlap. The proximal S coding sequence was isolated from pBH1-bam as a 1.5 kb Af1II-SphI fragment and ligated into Af1II-SphI digested pBH2 generating pCCV6. The full length S coding sequence was isolated as a 4.4 kb BamH1 fragment then ligated into the BamH1 site of the transfer vector RK19 to form pRKCCV6. Correct orientation of the gene was confirmed by restriction enzyme digestion. Thus, the plasmid RKCCV6 contains the CCV6 S gene downstream of the 4b promoter and flanked by TK coding sequences.

2.1.2. Isolation of recombinant virus

15

25

Recombinant vaccinia viruses were constructed by established procedures (Mackett & Smith, J.Gen.Virol. 67, 2067-2082, 1986). pRKCCV6 was transfected into vaccinia virus infected cells and recombinant viruses identified by dot-blot hybridisation using random primed ³²P labelled CCV6 spike gene as a probe Plaque purification and screening were repeated 3 times before stocks were prepared. The recombinant derived from pRKCCV6 was named Vac4b-C6.

2.2. Generation of vaccinia virus Vac4b-IN

2.2.1. Assembly of CCV Insavc-1 full length spike protein gene

The Insavc-1 (Bert) S gene was assembled from 3 overlapping cDNA clones BH8, BH9 and BH10. The cloning strategy is illustrated in figure 2. Digesting pBH8, which spans the middle of the S gene with Pvull and Hindlll yielded a 1.4 kb fragment. This fragment was ligated into a Pvull-Hindlll cut vector, pING14.2 forming pINGMS. This plasmid was linearized with Hindlll, phosphatased then gel eluted. The 3' S gene coding sequence isolated as a 1.1 kb Hindlll fragment from pBH10, was subcloned into Hindlll cut pINGMS generating pINGM3'S. Correct orientation of the cloned Hindlll fragment was confirmed by restriction enzyme digestion. Before the remaining coding sequence was excised from pBH9 a unique BamHI site was introduced 10bps upstream of the peplomer AUG start codon by site-directed mutagenesis (figure 2). The 5' coding sequence of the S gene was isolated as a 1.9 kb Pvull fragment and the remaining S gene coding sequence, which was isolated as a 2.5 kb Pvull partial-EcoRl fragment from pING3'S, were ligated in a two fragment ligation to BamHI-EcoRl digested pUC118. The complete S protein gene coding sequence was isolated as a 4.4 kb BamH1 fragment and subcloned into the BamH1 cut transfer vector pRK19, generating pRKINSAVC-1. Correct orientation of the gene was confirmed by restriction enzyme digestion. Thus the plasmid RKINSAVC-1 contains the CCV-INSAVC-1 S gene downstream of the vaccinia 4b promoter and flanked by TK coding sequences.

2.2.2. Isolation of recombinant vaccinia virus

Plasmid RKINSAVC-1 was used to introduce the S gene coding sequence into vaccinia virus by transfection and selection for TK⁻ recombinants was as described by Mackett and Smith, (1986, ibid). Recombinant virus isolates identified by dot blot hybridisation with a ³²P labelled CCV6 S DNA probe were subjected to three rounds of plaque purification and virus stocks prepared. The recombinant derived from RKINSAVC-1 was named Vac4b-IN.

2.3. Generation of vaccinia virus Vac4b-C54

2.3.1. Assembly of CCV C54 full length S protein gene

The C54 S gene coding sequence was assembled from the 3 overlapping clones pBH3, pBH4 and pBH11. A unique BamH1 site was created 10 bps upstream of the peplomer AUG start codon by site-directed mutagenesis in the proximal clone, pBH3 generating pBH3-bam (figure 3). A 2.0 kb Affill-EcoRI fragment was isolated from this plasmid and ligated to Affill-EcoRI digested pBH4 forming pBH5'MS. This plasmid was cleaved with Hindlll, phosphatased and gel eluted. The 3' coding sequence was excised as a

1.1 kb HindIII fragment from pBH11, then ligated to the HindIII digested pBH5'MS generating pBHC54. The correct orientation of the subcloned HindIII fragment was determined by restriction enzyme digestion. The full length C54 S gene was excised by digestion with BamH1 from pBHC54 and ligated into the BamH1 cut transfer vector RK19, forming pRKC54. Similarly, the orientation of the S gene was determined by restriction enzyme digestion. Thus the plasmid RKC54 contains the CCV C54 S gene downstream of the 4b promoter and flanked by TK coding sequences. The cloning strategy is illustrated in figure 3.

2.3.2. Isolation of recombinant vaccinia virus

Plasmid RKC54 was transfected into vaccinia virus infected cells. TK⁻ recombinants were selected using BUdR (Mackett and Smith, 1986, ibid). Recombinant virus isolates were identified by dot-blot hybridisation and subjected to three rounds of plaque purification before stocks were made. The recombinant derived from pRKC54 was named Vac4b-C54.

15 Example 3

Immunization experiments with live recombinant Vaccinia vaccine

3.1. Immunization

20

25

30

10

Cats were vaccinated with the following vaccines (107 pfu/cat):

- (a) 4 cats Vac4b-iN
- (b) 4 cats Vac4b-C6
- (c) 2 cats Vac4b-gB

(Vac4b-gB is recombinant Vaccinia virus which expresses the Cytomegalovirus glycoprotein gB under control of the 4b promoter)

(d) 2 cats - unvaccinated.

All cats were bled prior to vaccination (Bleed A). 3 weeks after vaccination the cats were bled again (Bleed B) and subsequently re-vaccinated as above.

2 weeks after re-vaccination all cats were bled (Bleed C).

3.2. Immuno-precipitation

Canine A72 cells were infected at a m.o.i. of about 10 with the recombinant viruses or mock-infected, incubated for 16 hours and starved of methionine for 1 hour. Infected cells were labelled with ³⁵S methionine and incubated for 30 min., washed and subsequently lysed in R.1.P.A. buffer. 1 µl cat antiserum (Bleed C) was added to the radiolabelled lysate and incubated on ice for 60 min. Protein G is added and incubated on ice for 60 min. After washing the protein G in R.1.P.A. buffer and PBS buffer, the bound proteins are recovered with 2% SDS 2% 2-mercapto-ethanol. The proteins are separated on 10% SDS polyacrylamide gel.

Sera from Bleed C precipitated the spike protein in the case of cats given Vac4b-C6 and Vac4b-IN. Thus, the cats immunized with the Vaccinia recombinant virus containing the spike genes responded with antibodies to the spike genes.

5 Legends to the Figures

- Figure 1: shows the cloning strategy for the construction of plasmid pRKCCV6 from plasmids pBH1 and pBH2.
- Fugure 2: shows the cloning strategy for the construction of plasmid pRKINSAVC-1 from plasmids pBH8, pBH10 and pBH9.
- Figure 3: shows the cloning strategy for the construction of plasmid pRKC54 from plasmids pBH3, pBH4 and pBH11.

55

SEQUENCE LISTING

5	(1) GENERAL INFORMATION:	
10	(i) APPLICANT: (A) NAME: AKZO N.V. (B) STREET: Velperweg 76 (C) CITY: Arnhem (E) COUNTRY: the Netherlands (F) POSTAL CODE (ZIP): 6824 BM	
	(ii) TITLE OF INVENTION: CANINE CORONAVIRUS SUBUNIT VACCINE	
	(iii) NUMBER OF SEQUENCES: 6	
15	 (v) COMPUTER READABLE FORM: (A) MEDIUM TYPE: Floppy disk (B) COMPUTER: IBM PC compatible (C) OPERATING SYSTEM: PC-DOS/MS-DOS (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 	
20	(vii) PRIOR APPLICATION DATA: (A) APPLICATION NUMBER: EP 91.303.737.0 (B) FILING DATE: 25-Apr-1991 (C) CLASSIFICATION:	
25	(2) INFORMATION FOR SEQ ID NO:1:	
30	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 4500 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear 	
	(ii) MOLECULE TYPE: cDNA	
35	(vi) ORIGINAL SOURCE:(A) ORGANISM: Canine corona virus(B) STRAIN: CCV-6	
10	<pre>(ix) FEATURE: (A) NAME/KEY: CDS (B) LOCATION: 654393 (D) OTHER INFORMATION: /label= CCV6_Spikegene</pre>	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:	
	GTAAGTTGCT CATTAGAAAC AATGGAAAAC TACTAAACTT TGGTAATCAC TTGGTTAATG	61
15	TGCC ATG ATT GTG CTA ATA TTG TGC CTC CTC TTG TTT TCG TAC AAT AGT Met Ile Val Leu Ile Leu Cys Leu Leu Phe Ser Tyr Asn Ser 1 5 10 15	.09
o		

12

		TG T Cys					_	_	_	_	_	1	.57
5		GGC Gly											05
10		GAA Glu 50		Val								2	253
		AAC Asn										3	101
15		ATA Ile							_			3	49
20	_	AAT Asn	_							_	_	3	97
		AGT Ser										4	45
25		TTG Leu 130										4	93
30		GCA Ala									_	5	41
		TCT Ser										5	89
35		AAT Asn									_	б	37
40		AAC Asn										6	85
		TCA Ser 210		_	_					_		7	733
45		GTT Val										7	781

50

5	TTG Leu 240	Lys	AGC Ser	TAT Tyr	GAA Glu	TTG Leu 245	Cys	GAA Glu	GAT Asp	TAT Tyr	GAA Glu 250	Tyr	TGC Cys	ACT	GGC Gly	TAT Tyr 255	829
3	GCT Ala	ACC Thr	AAC Asn	GTA Val	TTT Phe 260	Ala	CCG Pro	ACA Thr	GTG Val	GGC Gly 265	Gly	TAT Tyr	ATA Ile	CCT Pro	GAT Asp 270	Gly	877
10	TTC Phe	AGT Ser	TTT Phe	AAC Asn 275	AAT Asn	TGG Trp	TTT Phe	ATG Met	CTT Leu 280	ACA Thr	AAC Asn	AGT Ser	TCC Ser	ACG Thr 285	TTT Phe	GTT Val	925
15	AGT Ser	GGC Gly	AGA Arg 290	TTT Phe	GTA Val	ACA Thr	AAT Asn	CAA Gln 295	CCA Pro	TTA Leu	TTG Leu	GTT Val	AAT Asn 300	TGT Cys	TTG Leu	TGG Trp	973
	CCA Pro	GTG Val 305	CCC Pro	AGT Ser	TTT Phe	GGT Gly	GTC Val 310	GCA Ala	GCA Ala	CAA Gln	GAA Glu	TTT Phe 315	TGT Cys	TTT Phe	GAA Glu	GGT Gly	1021
20	GCG Ala 320	CAG Gln	TTT Phe	AGC Ser	CAA Gln	TGT Cys 325	AAT Asn	GGT Gly	GTG Val	TCT Ser	TTA Leu 330	AAC Asn	AAT Asn	ACA Thr	GTG Val	GAT Asp 335	1069
25	GTC Val	ATT Ile	AGA Arg	TTC Phe	AAC Asn 340	CTT Leu	AAT Asn	TTT Phe	ACC Thr	ACA Thr 345	GAT Asp	GTA Val	CAA Gln	TCT Ser	GGC Gly 350	ATG Met	1117
20	GGT Gly	GCT Ala	ATA Ile	GTA Val 355	TTT Phe	TCA Ser	CTG Leu	AAT Asn	ACA Thr 360	ACA Thr	GGT Gly	GGT Gly	GTC Val	ATT Ile 365	CTT Leu	GAG Glu	1165
30	ATT Ile	TCT Ser	TGT Cys 370	TAT Tyr	AAT Asn	GAT Asp	ACA Thr	GTG Val 375	AGT Ser	GAG Glu	TCA Ser	AGT Ser	TTC Phe 380	TAC Tyr	AGT Ser	TAT Tyr	1213
35	GGT Gly	GAA Glu 385	ATT	TCA Ser	ATC Ile	GGC Gly	GTA Val 390	ACT Thr	GAT Asp	GGA Gly	CCG Pro	CGT Arg 395	TAC Tyr	TGT Cys	TAC Tyr	GCC Ala	1261
	CTC Leu 400	TAT Tyr	AAT Asn	GGC Gly	CAG Gln	GCT Ala 405	CTT Leu	AAG Lys	TGT Cys	TTA Leu	GGA Gly 410	ACA Thr	TTA Leu	CCA Pro	CCT Pro	AGT Ser 415	1309
40	GTC Val	AAG Lys	GAA Glu	ATT Ile	GCT Ala 420	ATT Ile	AGT Ser	AAG Lys	TGG Trp	GGC Gly 425	CAT His	TTT Phe	TAT Tyr	ATT Ile	AAT Asn 430	GGT Gly	1357
45	TAC Tyr	AAT Asn	TTC Phe	TTT Phe 435	AGC Ser	ACT Thr	TTT Phe	CCT Pro	ATT Ile 440	GAT Asp	TGT Cys	ATA Ile	Ser	TTT Phe 445	AAT Asn	TTA Leu	1405
.~	ACC Thr	Thr	GGT Gly 450	GAT Asp	AGT Ser	GGA Gly	Ala	TTT Phe 455	TGG Trp	ACA Thr	ATT Ile	Ala	TAC Tyr 460	ACA Thr	TCG Ser	TAC Tyr	1453

50

														AAG Lys			1501
5		Cys												CTT Leu			1549
10						Phe							_	GTT Val	_		1597
														CAT His 525	_		1645
15														ATG Met			1693
20														GAT Asp			1741
														TAC Tyr			1789
25														TCC Ser			1837
30														ACT Thr 605			1885
														AAG Lys			1933
35	TTG Leu	TCA Ser 625	TTG Leu	AAT Asn	CCT Pro	GTT Val	GGT Gly 630	GCC Ala	AAC Asn	TGC Cys	AAG Lys	TTT Phe 635	GAT Asp	GTT Val	GCC Ala	GCT Ala	1981
40														GTA Val	_		2029
														AGT Ser			2077
45														TAT Tyr 685			2125

50

	TAT Tyr	GGT Gly	AGA Arg 690	Thr	GGT Gly	GTT Val	GGT Gly	ATT Ile 695	ATT Ile	AGA Arg	CAA Gln	ACT Thr	AAC Asn 700	AGT Ser	ACG Thr	CTA Leu	2173
5	CTT Leu	AGT Ser 705	GGC Gly	TTA Leu	TAT Tyr	TAC Tyr	ACA Thr 710	TCA Ser	CTA Leu	TCA Ser	GGT Gly	GAC Asp 715	TTG Leu	TTA Leu	GGG Gly	TTT Phe	2221
10	AAA Lys 720	AAT Asn	GTT Val	AGT Ser	Asp	GGT Gly 725	GTC Val	ATC Ile	ТАТ Туг	TCT Ser	GTC Val 730	ACG Thr	CCA Pro	TGT Cys	GAT Asp	GTA Val 735	2269
	AGC Ser	GTA Val	CAA Gln	GCT Ala	GCT Ala 740	GTT Val	ATT Ile	GAT Asp	GGC Gly	GCC Ala 745	ATA Ile	GTT Val	GGA Gly	GCT Ala	ATG Met 750	ACT Thr	2317
15	TCC Ser	ATT Ile	AAT Asn	AGT Ser 755	GAA Glu	CTG Leu	TTA Leu	GGT Gly	CTA Leu 760	ACA Thr	CAT His	TGG Trp	ACA Thr	ACA Thr 765	ACA Thr	CCT Pro	2365
20	AAT Asn	TTT Phe	TAT Tyr 770	TAT Tyr	TAT Tyr	TCT Ser	ATA Ile	TAT Tyr 775	AAT Asn	TAT Tyr	ACC Thr	AAT Asn	GAA Glu 780	AGG Arg	ACT Thr	CGT Arg	2413
	GGC	ACA Thr 785	GCA Ala	ATT Ile	GAT Asp	AGT Ser	AAC Asn 790	GAT Asp	GTT Val	GAT Asp	TGT Cys	GAA Glu 795	CCT Pro	ATC Ile	ATA Ile	ACC Thr	2461
25	TAT Tyr 800	TCT Ser	AAT Asn	ATA Ile	GGT Gly	GTT Val 805	TGT Cys	AAA Lys	AAT Asn	Gly	GCT Ala 810	TTG Leu	GTT Val	TTT Phe	ATT Ile	AAC Asn 815	2509
30	GTC Val	ACA Thr	CAT His	TCT Ser	GAT Asp 820	GGA Gly	GAC Asp	GTT Val	CAA Gln	CCA Pro 825	ATT Ile	AGC Ser	ACC Thr	GGT Gly	AAT Asn 830	GTC Val	2557
	ACG Thr	ATA Ile	CCT Pro	ACA Thr 835	AAT Asn	TTT Phe	ACC Thr	ATA Ile	TCT Ser 840	GTG Val	CAA Gln	GTT Val	GAA Glu	TAC Tyr 845	ATT Ile	CAG Gln	2605
35	GTT Val	TAC Tyr	ACT Thr 850	ACA Thr	CCG Pro	GTG Val	TCA Ser	ATA Ile 855	GAT Asp	TGT Cys	TCA Ser	AGG Arg	TAC Tyr 860	GTT Val	TGC Cys	AAT Asn	2653
40	GGT Gly	AAC Asn 865	CCT Pro	AGA Arg	TGC Cys	AAT Asn	AAA Lys 870	TTG Leu	TTA Leu	ACG Thr	CAA Gln	TAC Tyr 875	GTT Val	TCT Ser	GCA Ala	TGT Cys	2701
	CAA Gln 880	ACT Thr	ATT Ile	GAG Glu	CAA Gln	GCA Ala 885	CTT Leu	GCA Ala	ATG Met	GGT Gly	GCC Ala 890	AGA Arg	CTT Leu	GAA Glu	AAC Asn	ATG Met 895	2749
45	GAG Glu	ATT Ile	GAT Asp	TCC Ser	ATG Met 900	TTG Leu	TTT Phe	GTT Val	TCG Ser	GAA Glu 905	AAT Asn	GCC Ala	CTT Leu	AAA Lys	TTG Leu 910	GCA Ala	2797

50

	TCT Ser	GTT Val	GAA Glu	GCA Ala 915	TTA Leu	ATA Ile	GTA Val	GGA Gly	AAT Asn 920	TTA Leu	GAT Asp	CCT Pro	ATT Ile	TAC Tyr 925	AAA Lys	GAA Glu	2845
5	TGG Trp	CCT Pro	AAC Asn 930	ATT Ile	GGT Gly	GGT Gly	TCT Ser	TGG Trp 935	CTA Leu	GGA Gly	GGT Gly	TTA Leu	AAA Lys 940	GAC Asp	ATA Ile	TTG Leu	2893
10	CCA Pro	TCT Ser 945	CAC His	AAC Asn	AGC Ser	AAA Lys	CGT Arg 950	AAG Lys	TAC Tyr	CGG Arg	TCG Ser	GCT Ala 955	ATA Ile	GAA Glu	GAT Asp	TTG Leu	2941
	CTT Leu 960	TTT Phe	GAT Asp	AAG Lys	GTT Val	GTA Val 965	ACA Thr	TCT Ser	GGC Gly	TTA Leu	GGT Gly 970	ACA Thr	GTT Val	GAT Asp	GAA Glu	GAT Asp 975	2989
15	TAT Tyr	AAA Lys	CGT Arg	TGT Cys	ACA Thr 980	GGT Gly	GGT Gly	TAT Tyr	GAC Asp	ATA Ile 985	GCT Ala	GAC Asp	TTA Leu	GTG Val	TGT Cys 990	GCA Ala	3037
20	CAA Gln	TAT Tyr	TAC Tyr	AAT Asn 995	GGC Gly	ATC Ile	ATG Met	GTG Val	CTA Leu 1000	Pro	GGT Gly	GTA Val	GCT Ala	AAT Asn 1005	Asp	GAC Asp	3085
or-	AAG Lys	ATG Met	GCT Ala 1010	Met	TAC Tyr	ACT Thr	GCA Ala	TCT Ser 1015	Leu	GCA Ala	GGT Gly	GGT Gly	ATA Ile 1020	Thr	TTA Leu	GGT Gly	3133a
25	GCA Ala	CTT Leu 1025	Gly	GGT Gly	GGC Gly	GCA Ala	GTG Val 1030	Ser	ATA Ile	CCT Pro	TTT Phe	GCA Ala 1035	Ile	GCA Ala	GTT Val	CAA Gln	3181
30	GCC Ala 104	Arg	CTT Leu	AAT Asn	TAT Tyr	GTT Val 1045	Ala	CTA Leu	CAA Gln	ACT Thr	GAT Asp 1050	Val	TTG Leu	AAC Asn	AAG Lys	AAC Asn 1055	3229
	CAG Gln	CAG Gln	ATC Ile	CTG Leu	GCT Ala 1060	AAT Asn)	GCT Ala	TTC Phe	AAT Asn	CAA Gln 106	Ala	ATT Ile	GGT Gly	AAC Asn	ATT Ile 107	Thr	3277
35	CAG Gln	GCA Ala	TTT Phe	GGT Gly 1075	Lys	GTT Val	AAT Asn	GAT Asp	GCT Ala 1080	Ile	CAT His	CAA Gln	ACG Thr	TCA Ser 108	Gln	GGT Gly	3325
40	CTT Leu	GCT Ala	ACT Thr 109	Val	GCT Ala	AAA Lys	GCA Ala	TTG Leu 1095	Ala	AAA Lys	GTG Val	CAA Gln	GAT Asp 110	Val	GTT Val	AAC Asn	3373
	ACA Thr	CAA Gln 110	Gly	CAA Gln	GCT Ala	TTA Leu	AGC Ser 111	His	CTA Leu	ACA Thr	GTA Val	CAA Gln 111!	Leu	CAA Gln	AAT Asn	AAT Asn	3421
45		Gln				AGT Ser 1125	Ser					Tyr					3469

50 .

			AGT Ser			Ala					Leu					Leu	3517
5	ACA Thr	GCA Ala	CTT Leu	AAT Asn 115	Ala	TTT Phe	GTG Val	TCT Ser	CAG Gln 116	Thr	TTA Leu	ACC Thr	AGA Arg	CAA Gln 116	Ala	GAG Glu	3565
10	GTT Val	AGG Arg	GCT Ala 117	Ser	AGA Arg	CAG Gln	CTT Leu	GCT Ala 117	Lys	GAC Asp	AAG Lys	GTA Val	AAT Asn 1180	Glu	TGC Cys	GTT Val	3613
	AGG Arg	TCT Ser 118	CAA Gln 5	TCT Ser	CAG Gln	AGA Arg	TTT Phe 119	Gly	TTC Phe	TGT Cys	GGT Gly	AAT Asn 119	Gly	ACA Thr	CAT His	TTA Leu	3661
15	TTT Phe 120	Ser	CTT Leu	GCA Ala	AAT Asn	GCA Ala 120	Ala	CCA Pro	AAT Asn	GGC Gly	ATG Met 1210	Ile	TTC Phe	TTT Phe	CAC His	ACA Thr 1215	3709
20	GTG Val	CTA Leu	TTA Leu	CCA Pro	ACA Thr 1220	Ala	TAT Tyr	GAA Glu	ACC Thr	GTG Val 1225	Thr	GCC Ala	TGG Trp	TCA Ser	GGT Gly 1230	Ile	3757
	TGT Cys	GCA Ala	TCA Ser	GAT Asp 1235	Gly	GAT Asp	CGT Arg	ACT Thr	TTT Phe 1240	Gly	CTT Leu	GTT Val	GTT Val	AAG Lys 1245	Asp	GTC Val	3805
25	CAG Gln	TTG Leu	ACG Thr 1250	Leu	TTT Phe	CGC Arg	AAT Asn	CTA Leu 1255	Asp	GAC Asp	AAA Lys	TTC Phe	TAT Tyr 1260	Leu	ACT Thr	CCC Pro	3853
30	AGA Arg	ACT Thr 1265	ATG Met	TAT Tyr	CAG Gln	CCT Pro	AGA Arg 1270	Val	GCA Ala	ACT Thr	AGT Ser	TCT Ser 1275	Asp	TTT Phe	GTT Val	CAA Gln	3901
	ATT Ile 1280	Glu	GGA Gly	TGT Cys	Asp	GTG Val 1285	Leu	TTT Phe	GTT Val	Asn	GCA Ala 1290	Thr	GTA Val	ATT Ile	GAC Asp	TTG Leu 1295	3949
35	CCT Pro	AGT Ser	ATT Ile	ATA Ile	CCT Pro 1300	Asp	TAT Tyr	ATT Ile	GAT Asp	ATT Ile 1305	Asn	CAA Gln	ACT Thr	GTT Val	CAG Gln 1310	Asp	39 97
40	ATA Ile	TTA Leu	GAA Glu	AAT Asn 1315	Phe	AGA Arg	CCA Pro	AAT Asn	TGG Trp 1320	Thr	GTA Val	CCT Pro	GAG Glu	TTG Leu 1325	Pro	CTT Leu	4045
	GAC Asp	ATT Ile	TTC Phe 1330	Asn	GCA Ala	ACC Thr	TAC Tyr	TTA Leu 1335	Asn	CTG Leu	ACT Thr	GGT Gly	GAA Glu 1340	Ile	AAG Lys	TGC Cys	4093
45	TTA Leu	GAA Glu 1345	TTT Phe	AGG Arg	TCA Ser	Glu	AAG Lys 1350	Leu	CAT His	AAC Asn	Thr	ACA Thr 1355	Val	GAA Glu	CTT Leu	GCT Ala	4141

50

	ATT CTC ATT GAT AAT ATT AAT AAC ACA TTA TCA ATC TTA ATG CTC AAT Ile Leu Ile Asp Asn Ile Asn Asn Thr Leu Ser Ile Leu Met Leu Asn 1360 1365 1370 1375	4189
5	AGA ATT GAA ACT TAT GTA AAA TGG CCT TGG TAT GTG TGG CTA CTA ATT Arg Ile Glu Thr Tyr Val Lys Trp Pro Trp Tyr Val Trp Leu Leu Ile 1380 1385 1390	1237
10	GGA TTA GTA GTA ATA TTC TGC ATA CCC ATA TTG CTA TTT TGT TGT TGT Gly Leu Val Val Ile Phe Cys Ile Pro Ile Leu Leu Phe Cys Cys 1395	4285
	AGT ACT GGT TGT TGT GGA TGT ATT GGG TGT TTA GGA AGC TGT TGT CAT Ser Thr Gly Cys Cys Gly Cys Ile Gly Cys Leu Gly Ser Cys Cys His 1410 1415 1420	4333
15	TCC ATA TGT AGT AGA AGG CAA TTT GAA AGT TAT GAA CCA ATT GAA AAA Ser Ile Cys Ser Arg Arg Gln Phe Glu Ser Tyr Glu Pro Ile Glu Lys 1425 1430 1435	4381
20	GTT CAT GTT CAC TGAATTCAAA ATGTTAAGTC TACTATTTTA ATTACACCCG Val His Val His 1440	4433
~•	TGGCCACACA AGTTATATAA TGGTGCTGTC GTAAGTTCGA TACCAGTCAA CTATTAGCAT	4493
	TAATAAA	4500
25	(2) INFORMATION FOR SEQ ID NO:2:	
30	(i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 1443 amino acids (B) TYPE: amino acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear	
	(ii) MOLECULE TYPE: protein	
35	(vi) ORIGINAL SOURCE:(A) ORGANISM: Canina corona virus(B) STRAIN: CCV-6	·
40	<pre>(ix) FEATURE: (A) NAME/KEY: Protein (B) LOCATION: 11443 (D) OTHER INFORMATION: /label= CCV6_Spike</pre>	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:	
	Met Ile Val Leu Ile Leu Cys Leu Leu Phe Ser Tyr Asn Ser Val	

19

	Ile	Cys	Thr	Ser 20	Asn	Asn	Asp	Cys	Val 25	Gln	Val	Asn	Val	Thr 30	Gln	Leu
5	Pro	Gly	Asn 35	Glu	Asn	Ile	Ile	Lys 40	Asp	Phe	Leu	Phe	His 45	Thr	Phe	Lys
	Glu	Glu 50	Gly	Ser	Val	Val	Val 55	Gly	Gly	Tyr	Tyr	Pro 60	Thr	Glu	Val	Trp
10	Tyr 65	Asn	Cys	Ser	Arg	Ser 70	Ala	Thr	Thr	Thr	Ala 75	Tyr	Lys	Asp	Phe	Ser 80
	Asn	Ile	His	Ala	Phe 85	Tyr	Phe	Asp	Met	Glu 90	Asp	Met	Glu	Lys	Ser 95	Thr
15	Gly	Asn	Ala	Arg 100	Gly	Lys	Pro	Leu	Leu 105	Val	His	Val	His	Gly 110	Gly	Pro
	Val	Ser	Ile 115	Ile	Ile	Ile	Cys	Ala 120	Arg	Lys	Ala	Ser	Leu 125	Lys	His	Gly
20	Leu	Leu 130	Cys	Ile	Thr	Lys	Asn 135	Lys	Ile	Ile	Asp	Tyr 140	Asn	Thr	Phe	Thr
	Ser 145	λ la	Gln	Trp	Ser	Ala 150	Ile	Cys	Leu	Gly	Asp 155	Asp	Arg	Lys	Ile	Pro 160
25	Phe	Ser	Val	Ile	Pro 165	Thr	Asp	Asn	Gly	Thr 170	Lys	Ile	Phe	Gly	Leu 175	Glu
	Trp	Asn	Asp	Asp 180	Tyr	Val	Thr	Ala	Tyr 185	Ile	Ser	qaA	Arg	Ser 190	His	His
30	Leu	Asn	Ile 195	Asn	Asn	Asn	Trp	Phe 200	Asn	Asn	Val	Thr	Ile 205	Leu	Tyr	Ser
	Arg	Ser 210	Ser	Thr	Ala	Thr	Trp 215	Gln	Lys	Ser	Ala	Ala 220	Tyr	Val	Tyr	Gln
35	Gly 225	Val	Ser	Asn	Phe	Thr 230	Tyr	Tyr	Lys	Leu	Asn 235	Asn	Thr	Asn	Gly	Leu 240
	Lys	Ser	Tyr	Glu	Leu 245	Cys	Glu	Asp	Tyr	Glu 250	Tyr	Cys	Thr	Gly	Tyr 255	Ala
	Thr	Asn	Val	Phe 260	Ala	Pro	Thr	Val	Gly 265	Gly	Tyr	Ile	Pro	Asp 270	Gly	Phe
40	Ser	Phe	Asn 275	Asn	ттр	Phe	Met	Leu 280	Thr	Asn	Ser	Ser	Thr 285	Phe	Val	Ser
	Gly	Arg 290	Phe	Val	Thr	Asn	Gln 295	Pro	Leu	Leu	Val	Asn 300	Cys	Leu	Trp	Pro
45	Val 305	Pro	Ser	Phe	Gly	Val 310	Ala	Ala	Gln	Glu	Phe 315	Cys	Phe	Glu	Gly	Ala 320

50

55 .

	Gln	Phe	Ser	Gln	Cys 325	Asn	Gly	Val	Ser	Leu 330	Asn	Asn	Thr	Val	Asp 335	Val	
5	Ile	Arg	Phe	Asn 340	Leu	Asn	Phe	Thr	Thr 345	Asp	Val	Gln	Ser	Gly 350	Met	Gly	
	Ala	Ile	Val 355	Phe	Ser	Leu	Asn	Thr 360	Thr	Gly	Gly	Val	11e 365	Leu	Glu	Ile	
10	Ser	Cys 370	Tyr	λsn	Asp	Thr	Val 375	Ser	Glu	Ser	Ser	Phe 380	Tyr	Ser	Tyr	Gly	
	Glu 385	Ile	Ser	Ile	Gly	Val 390	Thr	Asp	Gly	Pro	Arg 395	Tyr	Cys	Tyr	Ala	Leu 400	
15	Tyr	Asn	Gly	Gln	Ala 405	Leu	Lys	Cys	Leu	Gly 410	Thr	Leu	Pro	Pro	Ser 415	Val	
20	Lys	Glu	Ile	Ala 420	Ile	Ser	Lys	Trp	Gly 425	His	Phe	Tyr	Ile	Asn 430	Gly	Tyr	
			435				Pro	440					445				
2 5	Thr	Gly 450	Asp	Ser	Gly	Ala	Phe 455	Trp	Thr	Ile	Ala	Tyr 460	Thr	Ser	Tyr	Thr	A, **
	Asp 465	Ala	Leu	Val	Gln	Val 470	Glu	Asn	Thr	Ala	Ile 475	Lys	Lys	Val	Thr	Tyr 480	
30	Cys	Asn	Ser	His	Ile 485	Asn	Asn	Ile	Lys	Cys 490	Ser	Gln	Leu	Thr	Ala 495	Asn	
	Leu	Gln	Asn	Gly 500	Phe	Tyr	Pro	Val	Ala 505	Ser	Ser	Glu	Val	Gly 510	Leu	Val	
35	Asn	Lys	Ser 515	Val	Val	Leu	Leu	Pro 520	Ser	Phe	Tyr	Ser	His 525	Thr	Ser	Val	
40	Asn	Ile 530	Thr	Ile	Asp	Leu	Gly 535	Met	Lys	Arg	Ser	Val 540	Met	Val	Thr	Ile	
40	Ala 545	Ser	Thr	Leu	Ser	Asn 550	Ile	Thr	Leu	Pro	Met 555	Gln	Asp	Asn	Asn	Thr 560	
4 5	Asp	Val	Tyr	Cys	Ile 565	Arg	Ser	Asn	Gln	Phe 570	Ser	Val	Tyr	Val	His 575	Ser	
	Thr	Cys	Lys	Ser 580	Ser	Leu	Trp	Asp	Asp 585	Val	Phe	Asn	Ser	Asp 590	Cys	Thr	
50	Asp	Val	Leu 595		Ala	Thr	Ala	Val 600	Ile	Lys	Thr	Gly	Thr 605	Cys	Pro	Phe	

	Ser	Phe 610	Asp	Lys	Leu	Asn	Asn 615	Tyr	Leu	Thr	Phe	Asn 620	Lys	Phe	Cys	Leu
5	Ser 625	Leu	Asn	Pro	Val	Gly 630	Ala	Asn	Cys	Lys	Phe 635	Asp	Val	Ala	Ala	Arg 640
	Thr	Arg	Thr	Asn	Glu 645	Gln	Val	Val	Arg	Ser 650	Leu	Tyr	Val	Ile	Tyr 655	Glu
10	Glu	Gly	Asp	Asn 660	Ile	Ala	Gly	Val	Pro 665	Ser	Asp	Asn	Ser	Gly 670	Leu	His
	Asp	Leu	Ser 675	Val	Leu	His	Leu	Asp 680	Ser	Суз	Thr	Asp	Tyr 685	Asn	Ile	Tyr
15	Gly	Arg 690	Thr	Gly	Val	Gly	Ile 695	Ile	Arg	Gln	Thr	Asn 700	Ser	Thr	Leu	Leu
	Ser 705	Gly	Leu	Tyr	Tyr	Thr 710	Ser	Leu	Ser	Gly	Asp 715	Leu	Leu	Gly	Phe	Lys 720
20	Asn	Val	Ser	Asp	Gly 725	Val	Ile	Tyr	Ser	Val 730	Thr	Pro	Cys	Asp	Val 735	Ser
	Val	Gln	Ala	Ala 740	Val	Ile	qzA	Gly	Ala 745	Ile	Val	Gly	Ala	Met 750	Thr	Ser
25	Ile	Asn	Ser 755	Glu	Leu	Leu	Gly	Leu 760	Thr	His	Trp	Thr	Thr 765	Thr	Pro	Asn
	Phe	Tyr 770	Tyr	Tyr	Ser	Ile	Tyr 775	Asn	Tyr	Thr	Asn .	Glu 780	Arģ	Thr	Arg	Gly
30	Thr 785	Ala	Ile	qaA	Ser	Asn 790	Asp	Val	qzA	Cys	Glu 795	Pro	Ile	Ile	Thr	Tyr 800
	Ser	Asn	Ile	Gly	Val 805	Cys	Lys	Asn	Gly	Ala 810	Leu	Val	Phe	Ile	Asn 815	Val
35	Thr	His	Ser	Asp 820	Gly	Asp	Val	Gln	Pro 825	Ile	Ser	Thr	Gly	Asn 830	Val	Thr
,	Ile	Pro	Thr 835	Asn	Phe	Thr	Ile	Ser 840	Val	Gln	Val	Glu	Tyr 845	Ile	Gln	Val
40	Tyr	Thr 850	Thr	Pro	Val	Ser	lle 855	Asp	Cys	Ser	Arg	Tyr 860	Val	Cys	Asn	Gly
	Asn 865	Pro	Arg	Cys	Asn	Lys 870	Leu	Leu	Thr	Gln	Tyr 875	Val	Ser	Ala	Cys	Gln 880
4 5	Thr	Ile	Glu	Gln	Ala 885	Leu	Ala	Met	Gly	Ala 890	Arg	Leu	Glu	Asn	Met 895	Glu
	Ile	Asp	Ser	Met 900	Leu	Phe	Val	Ser	Glu 905	Asn	Ala	Leu	Lys	Leu 910	Ala	Ser

50

	Val	Glu	Ala 915	Leu	Ile	Val	Gly	Asn 920	Leu	Asp	Pro	Ile	Tyr 925	Lys	Glu	Trp
5	Pro	Asn 930	Ile	Gly	Gly	Ser	Trp 935	Leu	Gly	Gly	Leu	Lys 940	Asp	Ile	Leu	Pro
	Ser 945	His	Asn	Ser	Lys	Arg 950	Lys	Tyr	Arg	Ser	Ala 955	Ile	Glu	Asp	Leu	Leu 960
10	Phe	Asp	Lys		Val 965	Thr	Ser	Gly	Leu	Gly 970	Thr	Val	Asp	Glu	Asp 975	Tyr
	Lys	Arg	Cys	Thr 980	Gly	Gly	Tyr	Asp	Ile 985	Ala	Asp	Leu	Val	Cys 990	Ala	Gln
15	Tyr	Tyr	Asn 995	Gly	Ile	Met	Val	Leu 1000		Gly	Val	Ala	Asn 1005		Asp	Lys
	Met	Ala 1010		Tyr	Thr	Ala	Ser 1015		Ala	Gly	Gly	Ile 1020		Leu	Gly	Ala
20	Leu 1025		Gly	Gly	Ala	Val 1030		Ile	Pro	Phe	Ala 1035		Ala	Val	Gln	Ala 1040
	Arg	Leu	Asn	Tyr	Val 1045		Leu	Gln	Thr	Asp 1050		Leu	Asn	Lys	Asn 1055	
25	Gln	Ile	Leu	Ala 1060		Ala	Phe	Asn	Gln 1065	Ala	Ile	Gly	Asn	Ile 1070		Gln
	Ala	Phe	Gly 1075		Val	Asn	Asp	Ala 1080		His	Gln	Thr	Ser 1085		Gly	Leu
30			1075 Val	5				1080 Ala)	His Val			1085 Val	5		
30	Ala	Thr 1090	1075 Val	Ala	Lys	Ala	Leu 1095 His	1080 Ala) Lys		Gln	Asp 1100 Leu	1085 Val)	Val	Asn	Thr
30	Ala Gln 1105	Thr 1090 Gly	Val Val Gln	Ala Ala	Lys Leu	Ala Ser 1110	Leu 1095 His	1080 Ala Leu	Lys Thr	Val	Gln Gln 1115 Tyr	Asp 1100 Leu	Val) Gln	Val Asn	Asn Asn	Thr Phe 1120 Glu
	Ala Gln 1105 Gln	Thr 1090 Gly S	Val Val Gln	Ala Ala Ser	Lys Leu Ser 1125	Ser	Leu 1095 His	1080 Ala Leu Ser	Lys Thr Asp	Val Val Ile 1130	Gln Gln 1115 Tyr	Asp 1100 Leu S	Val Oln Arg	Val Asn Leu	Asn Asn Asp 113:	Thr Phe 1120 Glu
	Ala Gln 1105 Gln Leu	Thr 1090 Gly Ala Ser	Val Cln Ile	Ala Ala Ser Asp 1140	Lys Leu Ser 1129 Ala	Ala Ser 1110 Ser Gln	Leu 1095 His Ile Val	1080 Ala Leu Ser Asp	Lys Thr Asp Arg 1145	Val Val Ile 1130	Gln Gln 1115 Tyr)	Asp 1100 Leu Asn Thr	Val Cln Arg	Val Asn Leu Arg 1156	Asn Asn Asp 113! Leu	Thr Phe 1120 Glu Thr
35	Ala Gln 1105 Gln Leu Ala	Thr 1090 Gly Ala Ser	Val Gln Ile Ala Asn 115:	Ala Ala Ser Asp 1146 Ala	Lys Leu Ser 1129 Ala Phe	Ser 1110 Ser Ser Gln Val	Leu 1095 His Ile Val	Ala Leu Ser Asp Gln 1160	Lys Thr Asp Arg 1145	Val Val Ile 1130 Leu Leu	Gln Gln 1115 Tyr) Ile	Asp 1100 Leu Asn Thr	Val Gln Arg Gly Gln 1169	Val Asn Leu Arg 1150	Asn Asp 113: Leu Glu	Thr Phe 1120 Glu Thr

50

55

	Ser	Leu	Ala	Asn	Ala 120		Pro	Asn	Gly	Met 121		Phe	Phe	His	Thr 121	
5	Leu	Leu	Pro	Thr 1220		Tyr	Glu	Thr	Val 122		Ala	Trp	Ser	Gly 123		Cys
	Ala	Ser	Asp 123		Asp	Arg	Thr	Phe 1240		Leu	Val	Val	Lys 124		Val	Gln
10	Leu	Thr 125		Phe	Arg	Asn	Leu 1255		Asp	Lys	Phe	Tyr 1260		Thr	Pro	Arg
	Thr 126	Met 5	Tyr	Gln	Pro	Arg 1270		Ala	Thr	Ser	Ser 1275		Phe	Val	Gln	Ile 1280
15	Glu	Gly	Cys	Asp	Val 1285		Phe	Val	Asn	Ala 1290		Val	Ile	Asp	Leu 1295	
	Ser	Ile	Ile	Pro 1300		Tyr	Ile	Asp	Ile 1305		Gln	Thr	Val	Gln 1310		Ile
20	Leu	Glu	Asn 1315	Phe	Arg	Pro	Asn	Trp 1320		Val	Pro	Glu	Leu 1325		Leu	Asp
	Ile	Phe 1330	Asn O	Ala	Thr	Tyr	Leu 1335		Leu	Thr	Gly	Glu 1340		Lys	Cys	Leu
25	Glu 134	Phe 5	Arg	Ser	Glu	Lys 1350		His	Asn	Thr	Thr 1355		Glu	Leu	Ala	Ile 1360
	Leu	Ile	Asp	Asn	Ile 1365		Asn	Thr	Leu	Ser 1370		Leu	Met	Leu	Asn 1375	
30	Ile	Glu	Thr	Tyr 1380	Val	Lys	Trp	Pro	Trp 1385		Val	Trp	Leu	Leu 1390		Gly
	Leu	Val	Val 1395	Ile	Phe	Cys	Ile	Pro 1400		Leu	Leu	Phe	Cys 1405		Cys	Ser
35	Thr	Gly 1410	Cys)	Cys	Gly	Cys	Ile 1415		Cys	Leu	Gly	Ser 1420		Cys	His	Ser
	Ile 142	Cys 5	Ser	Arg	Arg	Gln 1430		Glu	Ser	Tyr	Glu 1435		Ile	Glu	Lys	Val 1440
10	His	Val	His													
	(2) INFO	RMATI	ON F	OR S	EQ I	D NO	:3:									
15	(i)	(B) (C)	LEN TYF STR	CHA GTH: PE: n ANDE POLOG	442 ucle DNES	9 ba ic a S: s	se p cid ingl	airs	:							

50

		(11)	MOI	LECUI	E TY	PE:	CDNA	1									
5		(vi)	(2	A) OF	RGANI	OURCE ISM: I: CO	Cani			na vi	irus						
		(ix)	(<i>I</i>	3) LC	ME/I	ŒY: ON: INFO	60	4412 PION:	2 : /la	bel=	= CC1	/Ins#	vc-1	_spi	kege	ene	
10					•••	•											
		(xi)	SEÇ	OUENC	E DI	ESCRI	PTIC)N: 5	EQ 1	D NO):3:						
	TTGC	TCA'I	TA C	AAA	CAATO	G TA	AACI	'ACTA	AAC	TTTC	GTA	ATCA	CTTC	GT I	'AATO	TGCC	59
15	ATG Met	ATT Ile	GTG Val	CTT Leu	ACA Thr 5	TTG Leu	TGC Cys	CTT Leu	TTC Phe	TTG Leu 10	TTT Phe	TTG Leu	TAC Tyr	AGT Ser	AGT Ser 15	GTG Val	107
20	AGC Ser	TGT Cys	ACA Thr	TCA Ser 20	AAC Asn	AAT Asn	GAC Asp	TGT Cys	GTA Val 25	CAA Gln	GTT Val	AAT Asn	GTG Val	ACA Thr 30	CAA Gln	CTG Leu	155
	CCT Pro	GGC Gly	AAT Asn 35	GAA Glu	AAT Asn	ATT Ile	ATC Ile	AAA Lys 40	GAT Asp	TTT Phe	CTA Leu	TTT Phe	CAG Gln 45	AAC Asn	TTT Phe	AAA Lys	203
25	GAA Glu	GAA Glu 50	GGA Gly	AGT Ser	TTA Leu	GTT Val	GTT Val 55	GGT Gly	GGT Gly	TAT Tyr	TAC Tyr	CCC Pro 60	ACA Thr	GAG Glu	GTG Val	TGG Trp	251
30	TAT Tyr 65	AAC Asn	TGT Cys	TCC Ser	ACA Thr	ACT Thr 70	CAA Gln	CAA Gln	ACT Thr	ACC Thr	GCT Ala 75	TAT Tyr	AAG Lys	TAT Tyr	TTT Phe	AGT Ser 80	299
						TAT Tyr											347
35	GGC Gly	AAT Asn	GCA Ala	CGT Arg 100	GGT Gly	AAA Lys	CCT Pro	TTA Leu	CTA Leu 105	GTA Val	CAT His	GTT Val	CAT His	GGT Gly 110	TAA Asn	CCT Pro	395
40	GTT Val	AGT Ser	ATC Ile 115	ATT Ile	GTT Val	TAC Tyr	ATA Ile	TCA Ser 120	GCT Ala	TAT Tyr	AGA Arg	GAT Asp	GAT Asp 125	GTG Val	CAA Gln	TTT Phe	443
	AGG Arg	CCG Pro 130	CTT Leu	TTA Leu	AAG Lys	CAT His	GGT Gly 135	TTA Leu	TTG Leu	TGT Cys	ATA Ile	ACT Thr 140	AAA Lys	AAT Asn	GAC Asp	ACC Thr	491
4 5	GTT Val 145	Asp	TAT Tyr	AAT Asn	AGC Ser	TTT Phe 150	ACA Thr	ATT Ile	AAC Asn	CAA Gln	TGG Trp 155	CGA Arg	GAC Asp	ATA	TGT	TTG Leu 160	539

50

				AGA Arg													587
5				TTT Phe 180													635
10	ATT Ile	AGT Ser	GAT Asp 195	GAG Glu	TCT Ser	CAC His	CGT Arg	TTG Leu 200	AAT Asn	ATC Ile	TAA Asn	AAT Asn	AAT Asn 205	TGG Trp	TTT Phe	AAC Asn	683
				CTC Leu													731
15	AGT Ser 225	GCT Ala	GCA Ala	TAT Tyr	GTT Val	TAT Tyr 230	CAA Gln	GGT Gly	GTT Val	TCA Ser	AAT Asn 235	TTT Phe	ACT Thr	TAT Tyr	TAC Tyr	AAG Lys 240	779
20				ACC Thr													827
				ACT Thr 260													875
25				CCT Pro													923
30				ACT Thr													971
				TGC Cys													1019
35	GAA Glu	TTT Phe	TGT Cys	TTT Phe	GAA Glu 325	GGT Gly	GCT Ala	CAG Gln	TTT Phe	AGC Ser 330	CAA Gln	TGT Cys	AAC Asn	GGT Gly	GTT Val 335	TCT Ser	1067
40				ACA Thr 340													1115
				TCT Ser													1163
45	GGC	GGT Gly 370	GTC Val	ATT Ile	CTT Leu	GAG Glu	ATT Ile 375	TCT Ser	TGT Cys	TAT Tyr	AAT Asn	GAC Asp 380	ACA Thr	GTG Val	AGT Ser	GAG Glu	1211

50

						TAT Tyr 390											1259
5						GTA Val										Leu	1307
10						AGT Ser											1355
						GGT Gly											1403
75						TTA Leu											1451
20						TAC Tyr 470											1499
						ACG Thr											1547
25						GCT Ala											1595
30						CTT Leu											1643
						AGT Ser											1691
35						ACC Thr 550											1739
40	CCA Pro	ATG Met	CAG Gln	GAT Asp	AAT Asn 565	AAC Asn	ATA Ile	GAC Asp	GTG Val	TAC Tyr 570	TGT Cys	ATT Ile	CGT Arg	TCT Ser	AAC Asn 575	CAA Gln	1787
						CAT His											1835
<i>4</i> 5						TGT Cys											1883

50

55

	AAA Lys	ACT Thr 610	Gly	ACT Thr	TGT Cys	CCT Pro	TTC Phe 615	TCA Ser	TTT Phe	GAT Asp	AAA Lys	TTG Leu 620	AAT Asn	AAT Asn	TAC Tyr	TTA Leu	1931
5	ACT Thr 625	TTT Phe	AAC Asn	AAG Lys	TTC Phe	TG T Cys 630	TTG Leu	TCG Ser	TTG Leu	AAT Asn	CCC Pro 635	GTT Val	GGT Gly	GCC Ala	AAC Asn	TGT Cys 640	1979
10	AAG Lys	TTA Leu	GAT Asp	GTT Val	GCC Ala 645	GCC Ala	CGT Arg	ACA Thr	AGA Arg	ACC Thr 650	AAT Asn	GAG Glu	CAG Gln	GTT Val	TTT Phe 655	GGA Gly	2027
	AGT Ser	TTA Leu	TAT Tyr	GTA Val 660	ATA Ile	TAT Tyr	GAA Glu	GAA Glu	GGA Gly 665	GAC Asp	AAC Asn	ATA Ile	GTG Val	GGT Gly 670	GTA Val	CCG Pro	2075
15	TCT Ser	GAT Asp	AAT Asn 675	AGT Ser	GGT Gly	TTG Leu	CAC His	GAT Asp 680	TTG Leu	TCA Ser	GTG Val	TTG Leu	CAC His 685	TTA Leu	GAC Asp	TCT Ser	2123
20	TGT Cys	ACA Thr 690	GAT Asp	TAC Tyr	AAT Asn	ATA Ile	TAT Tyr 695	GGT Gly	AGA Arg	ACT Thr	GGT Gly	GTT Val 700	GGT Gly	ATT Ile	ATT Ile	AGA Arg	2171
	AAA Lys 705	ACT Thr	AAC Asn	AGC Ser	ACA Thr	CTA Leu 710	CTT Leu	AGT Ser	GGC Gly	TTA Leu	TAT Tyr 715	TAC Tyr	ACA Thr	TCA Ser	CTA Leu	TCA Ser 720	2219
25	GGT Gly	GAT Asp	TTG Leu	TTA Leu	GGT Gly 725	TTT Phe	AAA Lys	AAT Asn	GTT Val	AGT Ser 730	GAT Asp	GGT Gly	GTT Val	GTC Val	TAC Tyr 735	TCT Ser	2267
30	GTA Val	ACG Thr	CCA Pro	TGT Cys 740	GAT Asp	GTA Val	AGT Ser	GCA Ala	CAA Gln 745	GCT Ala	GCT Ala	GTT Val	ATT Ile	GAT Asp 750	GGT Gly	GCC Ala	2315
	ATA Ile	GTT Val	GGA Gly 755	GCT Ala	ATG Met	ACT Thr	TCC Ser	ATT Ile 760	AAT Asn	AGT Ser	GAA Glu	CTG Leu	TTA Leu 765	ggt Gly	CTA Leu	ACT Thr	2363
35	CAT His	TGG Trp 770	ACA Thr	ACA Thr	ACA Thr	CCT Pro	AAT Asn 775	TTT Phe	TAT Tyr	TAC Tyr	TAC Tyr	TCC Ser 780	ATA Ile	TAT Tyr	AAT Asn	TAT Tyr	2411
40	ACA Thr 785	AAT Asn	GTG Val	ATG Met	TAA Asn	CGT Arg 790	GGC Gly	ACG Thr	GCA Ala	ATT Ile	GAT Asp 795	AAT Asn	GAT Asp	ATT Ile	GAT Asp	TGT Cys 800	2459
	GAA Glu	CCT Pro	ATC Ile	ATA Ile	ACA Thr 805	TAT Tyr	TCT Ser	AAT Asn	ATA Ile	GGT Gly 810	GTT Val	TGT Cys	AAA Lys	AAT Asn	GGA Gly 815	GCT Ala	2507
45	TTG Leu	GTT Val	Phe	ATT Ile 820	AAC Asn	GTC Val	ACA Thr	CAT His	TCT Ser 825	GAT Asp	GGA Gly	GAC Asp	GTT Val	CAA Gln 830	CCA Pro	ATT Ile	2555

50

	ACC Thr		 -										2603
5	GAA Glu 850												2651
10	TAC Tyr		Asn										2699
45	GTT Val												2747
15	CTT Leu												2795
20	CTT Leu												2843
05	CCT Pro 930												2891
25	TTA Leu												2939
30	GCT Ala												2987
	ACA Thr											 	3035 ⁻
35	GAC Asp						Tyr				Val		3083
40	GTA Val 1010	Ala				Met				Ala			3131
	GGT Gly 5				Ala				Ala				3179
45	GCA Ala			Gln				Tyr				Thr	3227

50

55

. . .

					Lys	AAC Asn				Leu					Asn		3275
5	GCT Ala	ATT	GGT Gly 107	Asn	ATT Ile	ACA Thr	CAG Gln	GCA Ala 108	Phe	GGT Gly	AAG Lys	GTT Val	AAT Asn 108	Asp	GCT Ala	ATA Ile	3323
10	CAT His	CAA Gln 109	Thr	TCA Ser	AAA Lys	GGT Gly	CTT Leu 109	Ala	ACT Thr	GTT Val	GCT Ala	AAA Lys 1100	Ala	TTG Leu	GCA Ala	AAG Lys	3371
	GTG Val 1109	Gln	GAT Asp	GTT Val	GTT Val	AAC Asn 111	Thr	CAA Gln	GGT Gly	CAA Gln	GCT Ala 1115	Leu	AGC Ser	CAC His	CTA Leu	ACA Thr 1120	3419
15	GTA Val	CAA Gln	TTG Leu	CAA Gln	AAC Asn 112	AAT Asn õ	TTT Phe	CAA Gln	GCC Ala	ATT Ile 1130	Ser	AGT Ser	TCT Ser	ATT Ile	AGT Ser 113	Asp	3467
20	ATT Ile	TAT Tyr	AAC Asn	AGG Arg 1140	Leu	GAT Asp	GAA Glu	TTG Leu	AGT Ser 1145	Ala	GAT Asp	GCA Ala	CAA Gln	GTT Val 1150	Asp	AGG Arg	3515
	CTG Leu	ATT Ile	ACA Thr 1155	Gly	CGA Arg	CTT Leu	ACA Thr	GCA Ala 1160	Leu	AAT Asn	GCA Ala	Phe	GTG Val 1165	Ser	CAG Gln	ACT Thr	3563
25	TTA Leu	ACC Thr 1170	Arg	CAA Gln	GCA Ala	GAG Glu	GTT Val 1175	Arg	GCT Ala	AGT Ser	AGA Arg	CAA Gln 1180	Leu	GCT Ala	AAA Lys	GAC Asp	3611
30	AAG Lys 1185	Val	AAT Asn	GAA Glu	TGC Cys	GTT Val 1190	Arg	TCT Ser	CAA Gln	TCC Ser	CAG Gln 1195	Arg	TTT Phe	GGA Gly	TTC Phe	TGT Cys 1200	3659
	GGT Gly	AAT Asn	GGT Gly	ACA Thr	CAT His 1205	TTG Leu	TTT Phe	TCA Ser	CTT Leu	GCA Ala 1210	Asn	GCG Ala	GCA Ala	CCA Pro	AAT Asn 1215	Gly	3707
35	ATG Met	ATT Ile	TTC Phe	TTT Phe 1220	His	ACA Thr	GTG Val	CTA Leu	TTA Leu 1225	Pro	ACA Thr	GCT Ala	TAT Tyr	GAA Glu 1230	Thr	GTG Val	3755
40	ACG Thr	GCC [.] Ala	TGG Trp 1235	Ser	GGT Gly	ATT Ile	TGT Cys	GCG Ala 1240	Ser	GAT Asp	GGC Gly	AGT Ser	CGC Arg 1245	Thr	TTT Phe	GGA Gly	3803
	CTT Leu	GTT Val 1250	Val	GAG Glu	GAT Asp	GTC Val	CAG Gln 1255	Leu	ACG Thr	CTA Leu	TTT Phe	CGC Arg 1260	Asn	TTA Leu	GAT Asp	GAA Glu	3851
45	AAA Lys 1265	Phe	TAT Tyr	TTG Leu	ACG Thr	CCC Pro 1270	Arg	ACT Thr	ATG Met	TAT Tyr	CAG Gln 1275	Pro	AGA Arg	GTT Val	GCA Ala	ACT Thr 1280	3899

50

	AGT Ser	TCT Ser	GAT Asp	TTT Phe	GTT Val 1285	Gln	ATA Ile	GAA Glu	GGC	TGT Cys 1290	Asp	GTG Val	TTG Leu	TTT Phe	Val 1295	Asn	3947
5	GGA Gly	ACT Thr	GTA Val	ATT Ile 1300	Glu	TTG Leu	CCT Pro	AGT Ser	ATC Ile 1305	Ile	CC T Pro	GAC Asp	TAT Tyr	ATC Ile 1310	Asp	ATT Ile	3995
10	AAT Asn	CAA Gln	ACT Thr 1315	Val	CAG Gln	GAC Asp	ATA Ile	TTA Leu 1320	GAA Glu	AAT Asn	TTC Phe	AGA Arg	CCA Pro 1325	Asn	TGG Trp	ACT Thr	4043
	GTA Val	CCC Pro 1330	Glu	TTG Leu	CCA Pro	CTT Leu	GAC Asp 1335	Ile	TTT Phe	CAT His	GCA Ala	ACC Thr 1340	Tyr	TTA Leu	AAC Asn	CTG Leu	4091
15	ACT Thr 1345	Gly	GAA Glu	ATT	AAT Asn	GAC Asp 1350	Leu	GAA Glu	TTT Phe	AGG Arg	TCA Ser 1355	Glu	AAG Lys	TTA Leu	CAT His	AAC Asn 1360	4139
20	ACC Thr	ACA Thr	GTA Val	GAA Glu	CTT Leu 1365	Ala	ATT Ile	CTC Leu	ATT Ile	GAT Asp 1370	Asn	ATT Ile	AAT Asn	AAC Asn	ACA Thr 1375	Leu	4187
	GTC Val	AAT Asn	CTT Leu	GAA Glu 1380	Trp	CTC Leu	AAC Asn	AGA Arg	ATT Ile 1389	Glu	ACT Thr	TAT Tyr	GTA Val	AAA Lys 1390	\mathtt{Trp}	CCT Pro	4235
25	TGG Trp	TAT Tyr	GTT Val 1395	Trp	CTA Leu	CTA Leu	ATT Ile	GGA Gly 1400	TTA Leu)	GTA Val	GTA Val	ATA Ile	TTC Phe 140	Cys	ATA Ile	CCC Pro	4283
30	ATA Ile	TTG Leu 1410	Leu	TTT Phe	TGT Cys	TGT Cys	TGT Cys 1419	Ser	ACT Thr	GGT Gly	TGT Cys	TGT Cys 142	Gly	TGT Cys	ATC Ile	GJA GCG	4331
	TGT Cys 1425	Leu	GGA Gly	AGC Ser	TGT Cys	TGT Cys 1430	His	TCC Ser	ATA Ile	TGT Cys	AGT Ser 143	Arg	GGC Gly	CAA Gln	TTT Phe	GAA Glu 1440	4379
35	AGT Ser	TAT Tyr	GAA Glu	CCT Pro	ATT Ile 1449	Glu	AAA Lys	GTT Val	CAT His	GTT Val 1450	His	TGA	ATTC	AAA i	ATGT.	ГАА	4429

(2) INFORMATION FOR SEQ ID NO:4: 40

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1451 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

50

45

	(vi)) OR	GANI	SM:	: Cani: VInS			a vi	rus						
5	(ix)	(A (B) NAI	ME/KI	ON:	Prote lle RMAT	451	/lai	bel=	ccv	InSA	VC-1 _.	_Spi	ke		
10	(xi)	SEQ	UENC	E DES	SCRI	PTIO	N: S	EQ I	о ио	:4:						
	Met 1	Ile	Val	Leu	Thr 5	Leu	Cys	Leu	Phe	Leu 10	Phe	Leu	Tyr	Ser	Ser 15	Val
15	Ser	Cys	Thr	Ser 20	Asn	Asn	Asp	Cys	Val 25	Gln	Val	Asn	Val	Thr 30	Gln	Leu
	Pro	Gly	Asn 35	Glu	Asn	Ile	Ile	Lys 40	Asp	Phe	Leu	Phe	Gln 45	Asn	Phe	Lys
20	Glu	Glu 50	Gly	Ser	Leu	Val	Val 55	G] y	Gly	Туг	Tyr	Pro 60	Thr	Glu	Val	Trp
	Tyr 65	Asn	Cys	Ser	Thr	Thr 70	Gln	Gln	Thr	Thr	Ala 75	Tyr	Lys	Tyr	Phe	Ser 80
25	Asn	Ile	His	Ala	Phe 85	Tyr	Phe	Asp	Met	Glu 90	Ala	Met	Glu	Asn	Ser 95	Thr
	GJA	Asn	Ala	Arg 100	Gly	Lys	Pro	Leu	Leu 105	Val	His	Val	His	Gly 110	Asn	Pro
30	Val	Ser	Ile 115	Ile	Val	Tyr		Ser 120	Ala	Tyr	Arg	Asp	Asp 125	Val	Gln	Phe
	Arg	Pro 130	Leu	Leu	Lys	His	Gly 135	Leu	Leu	Cys	Ile	Thr 140	Lys	Asn	Asp	Thr
35	Val 145	Asp	Tyr	Asn	Ser	Phe 150	Thr	Ile	Asn	Gln	Trp 155	Arg	Asp	Ile	Cys	Leu 160
	Gly	Asp	Asp	Arg	Lys 165	Ile	Pro	Phe	Ser	Val 170	Val	Pro	Thr	Asp	Asn 175	Gly
40	Thr	Lys	Leu	Phe 180	Gly	Leu	Glu	Trp	Asn 185	Asp	Asp	Tyr	Val	Thr 190	Ala	Tyr
	Ile	Ser	Asp 195	Glu	Ser	His	Arg	Leu 200	Asn	Ile	Asn	Asn	Asn 205	Trp	Phe	Asn
1 5	Asn	Val 210	Thr	Leu	Leu	Tyr	Ser 215	Arg	Thr	Ser	Thr	Ala 220	Thr	Trp	Gln	His

32

50

	Ser 225	Ala	Ala	Tyr	Val	Туг 230	Gln	Gly	Val	Ser	Asn 235	Phe	Thr	Tyr	Tyr	Lys 240
5 .	Leu	Asn	Lys	Thr	Ala 245	Gly	Leu	Lys	Ser	Tyr 250	Glu	Leu	Cys	Glu	Asp 255	Tyr
	Glu	Tyr	Cys	Thr 260	Gly	Tyr	Ala	Thr	Asn 265	Val	Phe	Ala	Pro	Thr 270	Ser	Gly
10	Gly	Tyr	11e 275	Pro	Asp	Gly	Phe	Ser 280	Phe	Asn	Asn	Trp	Phe 285	Met	Leu	Thr
	Asn	Ser 290	Ser	Thr	Phe	Val	Ser 295	Gly	Arg	Phe	Val	Thr 300	Asn	Gln	Pro	Leu
15	Leu 305	Val	Asn	Cys	Leu	Trp 310	Pro	Val	Pro	Ser	Phe 315	Gly	Val	Ala	Ala	Gln 320
	Glu	Phe	Cys	Phe	Glu 325	Gly	Ala	Gln	Phe	Ser 330	Gln	Cys	Asn	Gly	Val 335	Ser
20	Leu	Asn	Asn	Thr 340	Val	Asp	Val	Ile	Arg 345	Phe	naA	Leu	Asn	Phe 350	Thr	Thr
	Asp	Val	Gln 355	Ser	Gly	Met	Gly	Ala 360	Thr	Val	Phe	Ser	Leu 365	Asn	Thr	Thr
25	Gly	Gly 370	Val	Ile	Leu	Glu	Ile 375	Ser	Cys	Tyr	Asn	Asp 380	Thr	Val	Ser	Glu
	Ser 385	Ser	Phe	Tyr	Ser	Tyr 390	Gly	Glu	Ile	Pro	Phe 395	Gly	Val	Thr	Asp	Gly 400
30	Pro	Arg	Tyr	Cys	Tyr 405	Val	Leu	Tyr	Asn	Gly 410	Thr	Ala	Leu	Lys	Tyr 415	Leu
	Gly	Thr	Leu	Pro 420	Pro	Ser	Val	Lys	Glu 425	Ile	Ala	Ile	Ser	Lys 430	Trp	Gly
35	His	Phe	Tyr 435	Ile	Asn	Gly	Tyr	Asn 440	Phe	Phe	Ser	Thr	Phe 445	Pro	Ile	Asp
	Cys	Ile 450	Ala	Phe	Asn	Leu	Thr 455	Thr	Gly	Ala	Ser	Gly 460	Ala	Phe	Trp	Thr
40	Ile 465	Ala	Tyr	Thr	Ser	Tyr 470	Thr	Glu	Ala	Leu	Val 475	Gln	Val	Glu	Asn	Thr 480
	Ala	Ile	Lys	Lys	Val 485	Thr	Tyr	Cys	Asn	Ser 490	His	Ile	Asn	Asn	Ile 495	Lys
45	Cys	Ser	Gln	Leu 500	Thr	Ala	Asn	Leu	Gln 505	Asn	Gly	Phe	Tyr	Pro 510	Val	Ala
	Ser	Ser	Glu 515	Val	Gly	Leu	Val	Asn 520	Lys	Ser	Val	Val	Leu 525	Leu	Pro	Ser

50

55

	Phe	Tyr 530	Ser	His	Thr	Ser	Val 535	Asn	Ile	Thr	Ile	Asp 540	Leu	Gly	Met	Lys
5	Arg 545	Ser	Val	Thr	Val	Thr 550	Ile	Ala	Ser	Pro	Leu 555	Ser	Asn	Ile	Thr	Leu 560
	Pro	Met	Gln	Asp	Asn 565	Asn	Ile	Asp	Val	Tyr 570	Cys	Ile	Arg	Ser	Asn 575	Gln
10	Phe	Ser	Val	Tyr 580	Val	His	Ser	Thr	Cys 585	Lys	Ser	Ser	Leu	Trp 590	Asp	Asn
	Asn	Phe	A≤n 595	Ser	Ala	Cys	Thr	Asp 600	Val	Leu	Asp	Ala	Thr 605	Ala	Va1	Ile
15	Lys	Thr 610	Gly	Thr	Cys	Pro	Phe 615	Ser	Phe	Asp	Lys	Leu 620	Asn	Asn	Tyr	Leu
	Thr 625	Phe	Asn	Lys	Phe	Cys 630	Leu	Ser	Leu	Asn	Pro 635	Val	Gly	Ala	Asn	Cys 640
20	Lys	Leu	Asp	Val	Ala 645	Ala	Arg	Thr	Arg	Thr 650	Asn	Glu	Gln	Val	Phe 655	Gly
	Ser	Leu	Tyr	Val 660	Ile	Tyr	Glu	Glu	Gly 665	qsA	Asn	Ile	Val	Gly 670	Val	Pro
25	Ser	Asp	Asn 675	Ser	Gly	Leu	His	Asp 680	Leu	Ser	Val	Leu	His 685	Leu	Asp	Ser
	Cys	Thr 690	Asp	Tyr	Asn	Ile	Tyr 695	Glÿ	Arg	Thr	Glý	Val 700	Gly	Ile	Ile	Arg
30	Lys 705	Thr	Asn	Ser	Thr	Leu 710	Leu	Ser	Gly	Leu	Tyr 715	Tyr	Thr	Ser	Leu	Ser 720
	Gly	Asp	Leu	Leu	Gly 725	Phe	Lys	Asn	Val	Ser 730	Asp	Gly	Val	Val	Tyr 735	Ser
35	Val	Thr	Pro	Cys 740	Asp	Val	Ser	Ala	Gln 745	Ala	Ala	Val	Ile	Asp 750	Gly	Ala
	Ile	Val	Gly 755	Ala	Met	Thr	Ser	Ile 760	Asn	Ser	Glu	Leu	Leu 765	Ğly	Leu	Thr
40	His	Trp 770	Thr	Thr	Thr	Pro	Asn 775	Phe	Tyr	Tyr	Tyr	Ser 780	Ile	Tyr	Asn	Tyr
	Thr 785	Asn	Val	Met	Asn	Arg 790	Gly	Thr	Ala	Ile	Asp 795	Asn	Asp	Ile	Asp	800 Cvs
45	Glu	Pro	Ile	Ile	Thr 805	Tyr	Ser	Asn	Ile	Gly 810	Val	Cys	Lys	Asn	Gly 815	Ala

50

	Leu	Val	Phe	Ile 820	Asn	Val	Thr	His	Ser 825	Asp	Gly	Asp	Val	Gln 830	Pro	Ile
5	Ser	Thr	Gly 835	Asn	Val	Thr	Ile	Pro 840	Thr	Asn	Phe	Thr	Ile 845	Ser	Val	Gln
	Val	Glu 850	Tyr	Ile	Gln	Val	Tyr 855	Thr	Thr	Pro	Val	Ser 860	Ile	Asp	Cys	Ala
	Arg 865	Tyr	Val	Cys	Asn	Gly 870	Asn	Pro	Arg	Cys	Asn 875	Lys	Leu	Leu	Thr	Gln 880
ı	Tyr	Val	Ser	Ala	Cys 885	Gln	Thr	Ile	Glu	Gln 890	Ala	Leu	Ala	Met	Gly 895	Ala
15	Arg	Leu	Glu	Asn 900	Met	Glu	Ile	Asp	Ser 905	Met	Leu	Phe	Val	Ser 910	Glu	Asn
	Ala	Leu	Lys 915	Leu	Ala	Ser	Val	Glu 920	Ala	Phe	Asn	Ser	Thr 925	Glu	Asn	Leu
20	Asp	Pro 930	Ile	Tyr	Lys	Glu	Trp 935	Pro	Asn	Ile	Gly	Gly 940	Ser	Trp	Leu	Gly
	Gly 945	Leu	Lys	Asp	Ile	Leu 950	Pro	Ser	His	Asn	Ser 955	Lys	Arg	Lys	Tyr	Arg 960
25	Ser	Ala	Ile	Glu	Asp 965	Leu	Leu	Phe	qaA	Lys 970	Val	Val	Thr	Ser	Gly 975	Leu
1	Gly	Thr	Val	Asp 980	Glu	Asp	Tyr	Lys	Arg 985	Ser	Ala	Gly	Gly	Tyr 990		Ile
30	Ala	Asp	Leu 995	Val	Cys	Ala	Arg	Tyr 1000		Asn	Gly	Ile	Met 1005		Leu	Pro
1	Gly	Val 1010		Asn	Asp	Asp	Lys 1015		Thr	Met	Tyr	Thr 1020		ser	Leu	Thr
	Gly 1025		Ile	Thr	Leu	Gly 1030		Leu	Ser	Gly	Gly 1035		Val	Ala	Ile	Pro 104
	Phe	Ala	Val	Ala	Val 1045		Ala	Arg	Leu	Asn 1050		Val	Ala	Leu	Gln 1055	
40	Asp	Val	Leu	Asn 1060		Asn	Gln	Gln	Ile 1069	Leu 5	Ala	Asn	Ala	Phe 1070		Gln
,	Ala	Ile	Gly 1075		Ile	Thr	Gln	Ala 1080		Gly	Lys	Val	Asn 1089		Ala	Ile
4 5	His	Gln 1090		Ser	Lys	Gly	Leu 1095		Thr	Val	Ala	Lys 1100		Leu	Ala	Lys
	Val 1105		Asp	Val	Val	Asn 1110		Gln	Gly	Gln	Ala 1115		Ser	His	Leu	Thr 112

35

50

	Val	Gln	Leu	Gln	Asn 112	Asn 5	Phe	Gln	Ala	Ile 113		Ser	Ser	Ile	Ser 113	
5	Ile	Tyr	Asn	Arg	Leu O	Asp	Glu	Leu	Ser 114		Asp	Ala	Gln	Val 115		Arg
	Leu	Ile	Thr 115		Arg	Leu	Thr	Ala 116		Asn	Ala	Phe	Val 116		Gln	Thr
10	Leu	Thr	Arg	Gln	Ala	Glu	Val 1175		Ala	ser	Arg	Gln 118		Ala	Lys	Asp
	Lys 118	Val	Asn	Glu	Cys	Val 1190	Arg)	Ser	Gln	Ser	Gln 1195		Phe	Gly	Phe	Cys 120
15	Gly	Asn	Gly	Thr	His 1205	Leu	Phe	Ser	Leu	Ala 1210		Ala	Ala	Pro	Asn 1215	
	Met	Ile	Phe	Phe 1220		Thr	Val	Leu	Leu 1225		Thr	Ala	Tyr	Glu 1230		Val
20	Thr	Ala	Trp 1235	Ser	Gly	Ile	Cys	Ala 1240		Asp	Gly	Ser	Arg 1245		Phe	Gly
	Leu	Val 1250	Val	Glu	Asp	Val	Gln 1255		Thr	Leu	Phe	Arg 1260		Leu	Asp	Glu
25	Lys 1265	Phe	Tyr	Leu	Thr	Pro 1270		Thr	Met	Tyr	Gln 1275		Arg	Val	Ala	Thr 128
•	Ser	Ser	Asp	Phe	Val 1285	Gln 5	Ile	Glu	Gly	Cys 1290		Val	Leu	Phe	Val 1295	
30	Gly	Thr	Val	Ile 1300	Glu)	Leu	Pro	Ser	Ile 1305		Pro	Ąsp	Tyr	Ile 1310		Ile
	Asn	Gln	Thr 1315	Val	Gln	Asp	Ile	Leu 1320		Asn	Phe	Arg	Pro 1325		Trp	Thr
35	Val	Pro 1330	Glu)	Leu	Pro	Leu	Asp 1335		Phe	His	Ala	Thr 1340		Leu	Asn	Leu
	Thr 1345	Gly	Glu	Ile	Asn	Asp 1350		Glu	Phe	Arg	Ser 1355		Lys	Leu	His	Asn 136
40	Thr	Thr	Val	Glu	Leu 1365	Ala	Ile	Leu	Ile	Asp 1370		Ile	Asn	Asn	Thr 1375	
	Val	Asn	Leu	Glu 1380	Trp	Leu	Asn	Arg	Ile 1385		Thr	Tyr	Val	Lys 1390		Pro
45	Trp	Tyr	Val 1395		Leu	Leu	Ile	Gly 1400		Val	Val	Ile	Phe 1405		Ile	Pro

50

		Ίle	Let 14:		Phe	Cys	Cys	Cys 141		Thị	Gly	y Cys	Cys 142	_	y Cys	Ile	Gly	
5		Cys 142		ı Gly	Ser	Cys	Cys 143		Ser	Ile	e Cys	s Ser 143		Gl)	7 Glr	Phe	Glu 144	
		Ser	Туг	r Glu	Pro	Ile 144		Lys	Val	. His	Val 145		i					
0	(2)	INFO	RMAT	NOI	FOR ·	SEQ	ID N	0:5:	•									
15		(i)	() (E	QUENC A) LE B) TY C) SI O) TO	NGTH PE: RAND	: 44 nucl EDNE	35 b eic SS:	ase acid sing	pair	s								
		(ii)	MOI	LECUI	E TY	PE:	cDNA											
20		(vi)	(P	GINA A) OR B) SI	GANI	SM:	Cani		oron	a vi	.rus							
		(ix)	(A	ATURE A) NA B) LO D) OT	ME/K	ON:	60			bel=	: CCV	7-C54	Spi	.kege	ne			
?5			•	•			٠		,	•				•				
		(xi)	SEC	QUENC	E DE	SCRI	PTIO	N: S	EQ I	D NO):5:							
_	TTG	CTCAT	TA G	SAAAC	AATG	G AA	AACT	ACTA	AAC	TTCC	GTA	ATCA	CTTG	GT I	'AATC	TGCC		59
30		ATT Ile																107
95		TGT Cys																155
		GGC Gly																203
10		GAA Glu 50	_			_			_									251
1 5		AAC Asn																299
50																		

37

	AAC Asn	ATA Ile	CAT His	GCA Ala	TTT Phe 85	Tyr	TTT Phe	GAT Asp	ATG Met	GAA Glu 90	GCT Ala	ATG Met	GCG Ala	AAT Asn	AGT Ser 95	ACT Thr	341
5	GGC	AAT Asn	GCA Ala	AGA Arg 100	GGT Gly	AAA Lys	CCT Pro	TTA Leu	CTA Leu 105	GTA Val	CAT His	GTT Val	CAT His	GGT Gly 110	AGT Ser	CCT Pro	395
10 .	GTT Val	AGT Ser	ATC Ile 115	ATT Ile	GTT Val	TAC Tyr	ATA Ile	TCA Ser 120	GCC Ala	TAT Tyr	AGA Arg	GAT Asp	GAT Asp 125	GTG Val	CAA Gln	AAT Asn	443
	AGG Arg	CCG Pro 130	CTC Leu	TTA Leu	AAA Lys	CAT His	GGT Gly 135	TTG Leu	TTG Leu	TGT Cys	ATA Ile	ACT Thr 140	AAA Lys	AAC Asn	AGC Ser	ACC Thr	491
15	ATT Ile 145	GAT Asp	TAT Tyr	AAC Asn	AGT Ser	TTT Phe 150	ACC Thr	TCT Ser	GCT Ala	CAG Gln	TGG Trp 155	CGT Arg	GAC Asp	ATA Ile	TGT Cys	TTG Leu 160	539
20	GGT Gly	ACT Thr	GAC Asp	AGA Arg	AAA Lys 165	ATA Ile	CCA Pro	TTC Phe	TCC Ser	GTC Val 170	GTA Val	ccc Pro	ACA Thr	GAT Asp	AAT Asn 175	GGC Gly	587
	ACA Thr	AAA Lys	CTA Leu	TTT Phe 180	GGT Gly	CTT Leu	GAG Glu	TGG Trp	ACT Thr 185	GAT Asp	GAC A sp	TAT Tyr	GTT Val	ACA Thr 190	GCC Ala	TAT Tyr	635
25	Ile	Ser	Asp 195	Asp	Ser	CAC His	Arg	Leu 200	Asn	Ile	Asn	Thr	Asn 205	Trp	Phe	Asn	683
30	AAT Asn	GTT Val 210	ACA Thr	ATC Ile	CTA Leu	TAC Tyr	TCC Ser 215	CGC Arg	TCA Ser	AGT Ser	Thr	GCC Ala 220	ACG Thr	TGG Trp	CAA Gln	AAG Lys	731
	AGT Ser 225	GCC Ala	GCA Ala	TAT Tyr	GTT Val	TAT Tyr 230	CAA Gln	GGT Gly	Val	Ser	AAT Asn 235	TTT Phe	ACG Thr	TAT Tyr	TAT Tyr	AAG Lys 240	779
35	Leu	Asn	Asn	Thr	Asn 245	GGC Gly	Leu	Lys	Ser	Tyr 250	Glu	Leu	Cys	Glu	Asp 255	Tyr	827
40	GAA Glu	TAC Tyr	TGC Cys	ACT Thr 260	GĞĈ Gly	TAT Tyr	GCC Ala	acc Thr	AAT Asn 265	GTG Val	TTT Phe	GCT Ala	CCG Pro	ACA Thr 270	TCA Ser	GGT Gly	875
	GGT Gly	Tyr	ATA Ile 275	CCT Pro	GAT Asp	GGA Gly	TTC Phe	AGT Ser 280	TTT Phe	AAC Asn	AAT Asn	TGG Trp	TTT Phe 285	ATG Met	CTT Leu	ACA Thr	923
45	Asn	AGC Ser 290	TCC Ser	ACT Thr	TTT Phe	GTT Val	AGT Ser 295	GGT Gly	AGG Arg	TTT Phe	GTA Val	ACA Thr 300	AAT Asn	CAA Gln	CCG Pro	CTG Leu	971

50

				TGC Cys													1019
5				TTT Phe													1067
70				ACA Thr 340		Asp											1115
	AAT Asn	GTA Val	CAA Gln 355	TCT Ser	GGC Gly	ATG Met	GGT Gly	GCT Ala 360	ACA Thr	GTA Val	TTT Phe	TCA Ser	CTG Leu 365	AAT Asn	ACA Thr	ACA Thr	1163
75				ATT Ile													1211
20				TAC Tyr													1259
				TGT Cys													1307
25				CCA Pro 420													1355
30	CAT His	TTT Phe	TAT Tyr 435	ATT Ile	TAA Asn	GGT Gly	TAC Tyr	AAT Asn 440	TTC Phe	TTT Phe	AGC Ser	ACT Thr	TTT Phe 445	CCT Pro	ATT Ile	GAT Asp	1403
				TTT Phe													1451
35				ACA Thr													1499
<i>4</i> 0				AAG Lys													1547
				CTT Leu 500													1595
45				GTT Val													1643

50

						AGT Ser											1691
5	CGT Arg 545	AGT Ser	GGT Gly	TAT Tyr	GGT Gly	CAA Gln 550	CCC Pro	ATA Ile	GCA Ala	TCA Ser	ACA Thr 555	CTA Leu	AGT Ser	AAC Asn	ATC Ile	ACA Thr 560	1739
10						AAT Asn											1787
						GTG Val											1835
15						GAC Asp											1883
20						TGT Cys											1931
						TTC Phe 630											1979
25	TGT Cys	AAG Lys	TTT Phe	GAT Asp	GTT Val 645	GCC Ala	GCC Ala	CGT Arg	ACA Thr	AGA Arg 650	ACC Thr	AAT Asn	GAG Glu	CAG Gln	GTT Val 655	GTT Val	2027
30	AGA Arg	AGT Ser	TTA Leu	TAT Tyr 660	GTA Val	ATG Met	TAT Tyr	GAA Glu	GAA Glu 665	GGA Gly	GAT Asp	AAC Asn	ATA Ile	GCG Ala 670	GGT Gly	GAC Asp	2075
						GGT Gly											2123
35						AAT Asn											2171
40						ACA Thr 710											2219
						GGT Gly											2267
45	TCT Ser	GTA Val	ACG Thr	CCA Pro 740	TGT Cys	GAT Asp	GTA Val	AGC Ser	GCA Ala 745	CAA Gln	GCT Ala	GCT Ala	GTT Val	ATT Ile 750	GAT Asp	GGT Gly	2315

50

	GCC Ala	ATA Ile	GTT Val 755	GGA Gly	GCT Ala	ATG Met	ACT Thr	TCC Ser 760	ATT Ile	AAT Asn	AGC Ser	GAA Glu	CTG Leu 765	TTA Leu	GGT Gly	CTA Leu	2363
5					ACA Thr												2411
10					AGA Arg												2459
					ATC Ile 805												2507
15					TTT Phe								Gly				2555
20					GGT Gly												2603
					TAC Tyr												2651
25	TGT Cys 865	GCA Ala	AGA Arg	TAC Tyr	GTT Val	TGC Cys 870	AAT Asn	GGT Gly	AAC Asn	CCT Pro	AGA Arg 875	TGC Cys	AAT Asn	AAA Lys	TTG Leu	TTA Leu 880	2699
30					TCT Ser 885												2747
	GGT Gly	GCC Ala	AGA Arg	CTT Leu 900	GAA Glu	AAC Asn	ATG Met	GAG Glu	ATT Ile 905	GAT Asp	TCC Ser	ATG Met	TTG Leu	TTT Phe 910	GTT Val	TCG Ser	2795
3 5					AAA Lys												2843
40	ACT Thr	CTA Leu 930	GAT Asp	CCT Pro	ATT Ile	TAC Tyr	AAA Lys 935	GAA Glu	TGG Trp	ccc Pro	AAT Asn	ATT Ile 940	GGT Gly	GGT Gly	TCT Ser	TGG Trp	2891
		Gly			AAA Lys						_						2939
45					ATA Ile 965												2987

50

55

	GGC Gly	TTA Leu	GGT Gly	ACA Thr 980	GTT Val	GAT Asp	GAA Glu	GAT Asp	TAT Tyr 985	AAG Lys	CGT Arg	TGT Cys	ACA Thr	GGT Gly 990	GGT Gly	TAT Tyr	3035
5				GAC Asp					Gln					Ile			3083
10	CTA Leu	Pro 101	Gly	GTA Val	GCT Ala	AAT Asn	GAT Asp 1015	Asp	AAG Lys	ATG Met	GCT Ala	ATG Met 1020	Tyr	ACT Thr	GCA Ala	TCT Ser	3131
	CTT Leu 102	Ala	GGT Gly	GGT Gly	ATA Ile	ACA Thr 1030	Leu	GGT Gly	GCA Ala	CTA Leu	GGT Gly 1035	Gly	GGC Gly	GCC Ala	GTG Val	GCT Ala 1040	3179
15	ATA Ile	CCT Pro	TTT Phe	GCA Ala	GTA Val 104	Ala	GTT Val	CAG Gln	GCT Ala	AGA Arg 1050	Leu	AAT Asn	TAT Tyr	GTT Val	GCT Ala 105	Leu	3227
20	CAA Gln	ACT Thr	GAT Asp	GTA Val 1060	Leu	AAC Asn	AAA Lys	AAC Asn	CAA Gln 1065	Gln	ATC Ile	CTG Leu	GCT Ala	AAT Asn 1070	Ala	TTC Phe	3275
	AAC Asn	CAA Gln	GCT Ala 1075	ATT Ile	GGT Gly	AAC Asn	ATT Ile	ACA Thr 1080	Gln	GCA Ala	TTT Phe	GGT Gly	AAG Lys 1089	Val	AAT Asn	GAC Asp	3323
25	GCA Ala	ATA Ile 1090	His	CAA Gln	ACA Thr	TCA Ser	CAA Gln 1095	Gly	CTT Leu	GCC Ala	ACT Thr	GTT Val 1100	Ala	AAA Lys	GCA Ala	TTG Leu	3371
30	GCA Ala 1105	Lys	GTG Val	CAA Gln	GAT Asp	GTT Val 1110	Val	AAC Asn	ACA Thr	CAA Gln	GGT Gly 1115	Gln	GCT Ala	TTA Leu	AGC Ser	CAC His 1120	3419
	CTA Leu	ACA Thr	GTA Val	CAA Gln	TTG Leu 1125	Gln	AAC Asn	AAT Asn	TTT Phe	CAA Gln 1130	Ala	ATT Ile	AGT Ser	AGT Ser	TCC Ser 1135	Ile	3467
35				TAC Tyr 1140	Asn					Leu					Gln		3515
40	GAC Asp	AGG Arg	CTT Leu 1155	ATT Ile	ACA Thr	GGA Gly	AGA Arg	CTT Leu 1160	Thr	GCA Ala	CTT Leu	AAT Asn	GCA Ala 1165	Phe	GTG Val	TCT Ser	3563
	Gln	ACT Thr 1170	Leu	ACC Thr	AGA Arg	CAA Gln	GCA Ala 1175	Glu	GTT Val	AGG Arg	GCT Ala	AGT Ser 1180	Arg	CAA Gln	CTT Leu	GCT Ala	3611
45	AAA Lys 1185	Asp	AAA Lys	GTT Val	AAT Asn	GAA Glu 1190	Cys	GTT Val	AGG Arg	TCT Ser	CAA Gln 1195	Ser	CAG Gln	AGA Arg	TTT Phe	GGA Gly 1200	3659

	TTC Phe	TG T Cys	GGT Gly	TAA Asn	GGT Gly 1205	Thr	CAT His	TTG Leu	TTT Phe	TCA Ser 1210	Leu	GCA Ala	AAT Asn	GCA Ala	GCA Ala 1215	Pro	3707
5	AAT Asn	GGC Gly	ATG Met	ATT Ile 1220	Phe	TTT Phe	CAC His	Thr	GTG Val 1225	Leu	TTA Leu	CCA Pro	ACA Thr	GCT Ala 1230	Tyr	GAA Glu	3755
10	ACT Thr	GTG Val	ACG Thr 1235	Ala	TGG Trp	TCA Ser	GGT Gly	ATT Ile 1240	Cys	GCA Ala	TCA Ser	GAT Asp	GGC Gly 1245	Asp	CGC Arg	ACT Thr	3803
	TTT Phe	GGA Gly 1250	Leu	GTT Val	GTT Val	AAG Lys	GAT Asp 1255	Val	CAG Gln	CTG Leu	ACG Thr	CTA Leu 1260	Phe	CGC Arg	AAT Asn	TTA Leu	3851
15	GAT Asp 1265	qzA	AAA Lys	TTC Phe	TAT Tyr	TTG Leu 1270	ACT Thr	CCC Pro	AGA Arg	ACT Thr	ATG Met 1275	Tyr	CAG Gln	CCT Pro	AGA Arg	GTT Val 1280	3899
20	GCA Ala	ACT Thr	AGT Ser	TCT Ser	GAT Asp 1285	Phe	GTT Val	CAA Gln	ATA Ile	GAA Glu 1290	Gly	TGT Cys	GAT Asp	GTG Val	TTG Leu 1295	Phe	3947
	GTC Val	AAT Asn	GCA Ala	ACT Thr 1300	Val	ATT Ile	GAC Asp	TTG Leu	CCT Pro 130	Ser	ATC Ile	ATA Ile	CCT Pro	GAC Asp 1310	Tyr	ATT Ile	3995
25				Gln			CAG Gln		Ile					Arg			4043
30			Val				ACA Thr 1335	Leu					Ala				4091
	AAC Asn 1349	Leu	ACT Thr	GGT Gly	GAA Glu	ATT Ile 1350	AAT Asn)	GAC Asp	TTA Leu	GAA Glu	TTT Phe 135	Arg	TCG Ser	GAA Glu	AAG Lys	TTA Leu 1360	4139
35	CAT His	AAC Asn	ACC Thr	ACA Thr	GTA Val 136	Glu	CTT Leu	GCT Ala	GTT Val	CTC Leu 1370	Ile	GAT Asp	AAT Asn	ATT Ile	AAT Asn 137	Asn	4187
40	ACA Thr	TTA Leu	GTC Val	AAT Asn 138	Leu	GAA Glu	TGG Trp	CTC Leu	AAT Asn 138	Arg	ATT Ile	GAA Glu	ACT Thr	TAT Tyr 139	٧al	AAA Lys	4235
	TGG Trp	CCT Pro	TGG Trp 139	Tyr	GTG Val	TGG Trp	CTA Leu	CTA Leu 140	Ile	GGA Gly	TTA Leu	GTA Val	GTA Val 140	Ile	TTC Phe	TGC Cys	4283
45							TGC		TGT Cys						_		4331

	ATA GGT Ile Gly 1425	TGC Cys	TTA Leu	GGA Gly	AGT Ser 1430	Cys	TGT Cys	CAC His	TCT Ser	ATG Met 1435	Cys	AGT Ser	AGA Arg	AGA Arg	CAA Gln 1440	•	4379
5	TTT GAA Phe Glu	AGT Ser	TAT Tyr	GAA Glu 1445	Pro	ACC Thr	GAA Glu	AAA Lys	GTG Val 1450	His	GTC Val	CAC His	TAAF	ATTCA	LAA		4428
	АСТААТА																4435
10																	
15	(2) INFO	SEQI (A (B	UENC } LE } TY } ST		ARAC : 14 amin EDNE	TERI 53 a o ac SS:	STIC mino id sing	S: aci	ds								
	(ii)	MOL	ECUL	E TY	PE:	prot	ein										
20	(vi)		OR	L SO GANI: RAIN	SM:	Cani		oron	a vi	rus							
25	(ix)	(B)	NAI LO	: ME/KI CATIO HER I	ON:	11	453	/la	bel=	ccv-	-C54 _.	_spi	ke				
	(xi)	SEQU	JENC:	e des	SCRI	PTIO	N: S	EQ I	D NO	:6:							
30	Met 1	Ile	Val	Leu	Thr 5	Leu	Cys	Leu	Leu	Leu 10	Phe	Ser	Tyr	Asn	Ser 15	۷al	
	Ile	Cys	Thr	Ser 20	Asn	Asn	Asp	Cys	Val 25	Gln	Val	Asn	Val	Thr 30	Gln	Leu	
35	Pro	Gly	Asn 35	Gļu	Asn	Ile	Ile	Lys 40	Asp	Phe	Leu	Phe	Gln 45	Asn	Phe	Lys	
	Glu	Glu 50	Gly	Ser	Val	Val	Val 55	Gly	Gly	Tyr	туг	Pro 60	Thr	Glu	Val.	Trp	
40	Tyr 65	Asn	Cys	Ser	Arg	Thr 70	Ala	Thr	Thr	Thr	Ala 75	Tyr	His	Tyr	Phe	Ser 80	
	Asn	Ile	His	Ala	Phe 85	Tyr	Phe	Asp	Met	Glu 90	Ala	Met	Ala	Asn	Ser 95	Thr	
45	Gly	Asn	Ala	Arg 100	Gly	Lys	Pro	Leu	Leu 105	Val	His	Val	His	Gly 110	Ser	Pro	
50									-								
			•														

	Val	Ser	Ile 115	Ile	Val	Tyr	Ile	Ser 120	Ala	Тух	Arg	Asp	Asp 125	Val	Gln	Asn
5	Arg	Pro 130	Leu	Leu	Lys	His	Gly 135	Leu	Leu	Cys	Ile	Thr 140	Lys	Asn	Ser	Thr
	11e 145	Asp	Tyr	Asn	Ser	Phe 150	Thr	Ser	Ala	Gln	Trp 155	Arg	Asp	Ile	Cys	Leu 160
10	Gly	Thr	Asp	Arg	Lys 165	Ile	Pro	Phe	Ser	Val 170	Val	Pro	Thr	Asp	Asn 175	Gly
	Thr	Lys	Leu	Phe 180	Gly	Leu	Glu	Trp	Thr 185	Asp	Asp	Tyr	Val	Thr 190	Ala	Tyr
15	Ile	Ser	Asp 195	Asp	Ser	His	Arg	Leu 200	Asn	Ile	Asn	Thr	Asn 205	Trp	Phe	Asn
	Asn	Val 210	Thr	Ile	Leu	Tyr	Ser 215	Arg	Ser	Ser	Thr	Ala 220	Thr	Trp	Gln	Lys
20	Ser 225	Ala	Ala	Tyr	Val	Tyr 230	Gln	Gly	Val	Ser	Asn 235	Phe	Thr	Tyr	Tyr	Lys 240
	Leu	Asn	Asn	Thr	Asn 245	Gly	Leu	Lys	Ser	Tyr 250	Glu	Leu	Cys	Glu	Asp 255	Tyr
25	Glu	Tyr	Cys	Thr 260	Gly	Tyr	Ala	Thr	Asn 265	Val	Phe	Ala	Pro	Thr 270	Ser	Gly
	Gly	Tyr	Ile 275	Pro	Asp	Gly	Phe	Ser 280	Phe	Asn	Asn	Trp	Phe 285	Met	Leu	Thr
30	Asn	Ser 290	Ser	Thr	Phe	Val	Ser 295	Gly	Arg	Phe	Val	Thr 300	Asn	Gln	Pro	Leu
	Leu 305	Val	Asn	Cys	Leu	Val 310	Pro	Val	Pro	Ser	Phe 315	Gly	Val	Ala	Ala	Gln 320
35	Glu	Phe	Cys	Phe	Glu 325	Gly	Ala	Gln	Phe	Ser 330	Gln	Cys	Asn	Gly	Val 335	Ser
	Leu	Asn	Asn	Thr 340	Val	Asp	Val	Ile	Arg 345	Phe	Asn	Leu	Asn	Phe 350	Thr	Thr
40	Asn	Val	Gln 355	Ser	Gly	Met	Gly	Ala 360	Thr	Val	Phe	Ser	Leu 365	Asn	Thr	Thr
	Gly	Gly 370	Val	Ile	Leu	Glu	Ile 375	Ser	Cys	Tyr	Asn	Asp 086	Thr	Val	Ser	Glu
4 5	Ser 385	Ser	Phe	Tyr	Ser	Tyr 390	Gly	Glu	Ile	Pro	Phe 395	Gly	Val	Thr	Asp	Gly 400
	Pro	Arg	Tyr	Суѕ	Tyr 405	Val	Leu	Tyr	naA	Gly 410	Thr	Ala	Leu	Lys	Tyr 415	Leu

45

	Gly	Thr	Leu	Pro 420	Pro	Ser	Val	Lys	Glu 425	Ile	Ala	Ile	ser	Lys 430		Gly
5	His	Phe	Tyr 435	Ile	Asn	Gly	Tyr	Asn 440		Phe	Ser	Thr	Phe 445		Ile	Asp
	Cys	Ile 450	Ser	Phe	Asn	Leu	Thr 455	Thr	Gly	Asp	Ser	Gly 460		Phe	Trp	Thr
10	Ile 465	Ala	Tyr	Thr	Ser	Tyr 470	Thr	Glu	Ala	Leu	Val 475	Gln	Val	Glu	Asn	Thr 480
	Ala	Ile	Lys	Lys	Val 485	Thr	Tyr	Cys	Asn	Ser 490	His	Ile	Asn	Asn	Ile 495	Lys
15	Cys	Ser	Gln	Leu 500	Thr	Ala	Asn	Leu	Gln 505	Asn	Gly	Phe	Tyr	Pro 510	Val	Ala
	Ser	Ser	Glu 515	Val	Gly	Leu	Val	Asn 520	Lys	Ser	Val	Val	Leu 525	Leu	Pro	Ser
20	Phe	Tyr 530	Ser	His	Thr	Ser	Val 535	Asn	Ile	Thr	Ile	Asp 540	Leu	Gly	Met	Lys
	Arg 545	Ser	Gly	Tyr	Gly	Gln 550	Pro	Ile	Ala	Ser	Thr 555	Leu	Ser	Asn	Ile	Thr 560
25	Leu	Pro	Met	Gln	Asp 565	Asn	Asn	Thr	Asp	Val 570	Tyr	Cys	Ile	Arg	Ser 575	Asn
	Gln	Phe	Ser	Val 580	Tyr	Val	His	Ser	Thr 585	Cys	Lys	Ser	Ser	Leu 590	Trp	Asp
30	Asn	Ile	Phe 595	Asn	Ser	Asp	Cys	Thr 600	Asp	Val	Leu	His	Ala 605	Thr	Ala	Val
	Ile	Lys 610	Thr	Gly	Thr	Cys	Pro 615	Phe	Ser	Phe	Asp	Lys 620	Leu	Asn	Asn	Tyr
35	Leu 625	Thr	Phë	Asn	Lys	Phe ⁻	Cys	Leu'	Ser	Leu	Asn 635	Pro	Val	Gly	Ala	Asn 640
	Cys	Lys	Phe	Asp	Val 645	Ala	Ala	Arg	Thr	Arg 650	Thr	Asn	Glu	Gln	Val 655	Val.
40	Arg	Ser	Leu	Tyr 660	Val	Met	Tyr	Glu	Glu 665	Gly	Asp	Asn	Ile	Ala 670	Gly	Asp
-	Arg	Pro	Asp 675	Asn	Ser	Gly	Leu	His 680	Asp	Leu	Ser	Val	Leu 685	His	Leu	Asp
45	Ser	Cys 690	Thr	Asp '	Tyr	Asn	Ile 695	Tyr	Gly	Arg		Gly 700	Val	Gly	Ile	Ile

50

	Arg 705	Gln	Thr	Asn	Ser	Thr 710	Ile	Phe	Ser	Gly	Leu 715	Tyr	Tyr	Thr	Ser	Leu 720
5	Ser	Gly	Asp	Leu	Leu 725	Gly	Phe	Lys	Asn	Val 730	Ser	Asp	Gly	Val	Val 735	Tyr
	Ser	Val	Thr	Pro 740	Cys	Asp	Val	Ser	Ala 745	Gln	Ala	Ala	Val	Ile 750	Asp	Gly
10	Ala	Ile	Val 755	Gly	Ala	Met	Thr	Ser 760	Ile	Asn	Ser	Glu	Leu 765	Leu	Gly	Leu
	Thr	His 770	Trp		Thr	Thr	Pro 775	Asn	Phe	Tyr	Tyr	Tyr 780	Ser	Ile	Tyr	Asn
15	Tyr 785	Thr	Ser	Val	Arg	Thr 790	Arg	Gl.y	Thr	Ala	Ile 795	Asp	Ser	Asn	Asp	Val 800
	Ąsp	Суѕ	Glu	Pro	Ile 805	Ile	Thr	Tyr	Ser	Asn 810	Ile	Gly	Val	Cys	Lys 815	Asn
20	Gly	Ala	Leu	Val 820	Phe	Ile	Asn	Val	Thr 825	His	Ser	Asp	Gly	Asp 830	Val	Gln
	Pro	Ile	Ser 835	Thr	Gly	Asn	Val	Thr 840	Ile	Pro	Thr	Asn	Phe 845	Thr	Ile	ser
25	Val	Gln 850	Val	Glu	Tyr	Ile	Gln 855	Val	Tyr	Thr	Thr	Pro 860	Val	Ser	Ile	Asp
	Cys 865	Ala	Arg	Tyr	Val	Cys 870	Asn	Gly	Asn	Pro	Arg 875	Cys	Asn	Lys	Leu	Leu 880
30	Thr	Gln	Tyr	Val	Ser 885	Ala	Cys	Gln	Thr	Ile 890	Glu	Gln	Ala	Leu	Ala 895	Met
	Gly	Ala	Arg	Leu 900	Glu	Asn	Met	Glu	11e 905	Asp	Ser	Met	Leu	Phe 910	Val	Ser
35	Glu	Asn	Ala 915	Leu	Lys	Leu	Ala	Ser 920	Val	Glu	Ala	Phe	Asn 925	Ser	Thr	Glu
	Thr	Leu 930	Asp	Pro	Ile	Tyr	Lys 935	Glu	Trp	Pro	Asn	Ile 940	Gly	Gly	Ser	Trp
40	Leu 945	Gly	Gly	Leu	Lys	Asp 950	Ile	Leu	Pro	Ser	His 955	Asn	Ser	Lys	Arg	Lys 960
	Tyr	Arg	Ser	Ala	11e 965	Glu	Asp	Leu	Leu	Phe 970	Asp	Lys	Val	Val	Thr 975	Ser
45	Gly	Leu	Gly	Thr 980	Val	Asp	Glu	Asp	Tyr 985	Lys	Arg	Cys	Thr	Gly 990	Gly	Tyr
	Asp	Ile	Ala 995	Asp	Leu	Val	Суз	Ala 1000		Tyr	Tyr	Asn	Gly 1005		Met	Val

47

50

	Leu	Pro 101	Gly o	Val	Ala	Asn	Asp 101	Asp 5	Lys	Met	Ala	Met 102		Thr	Ala	Ser
5	Leu 102	Ala 5	Gly	Gly	Ile	Thr 103	Leu 0	Gly	Ala	Leu	Gly 103		Gly	Ala	Val	Ala 1040
	Ile	Pro	Phe	Ala	Val 104	Ala 5	Val	Gln	Ala	Arg 105	Leu 0	Asn	Tyr	Val	Ala 105	
10	Gln	Thr	Asp	Val 106	Leu O	Asn	Lys	Asn	Gln 106		Ile	Leu	Ala	Asn 107		Phe
	Asn	Gln	Ala 107	Ile 5	Gly	Asn	Ile	Thr 1080	Gln O	Ala	Phe	Gly	Lys 108	Val	Asn	Asp
15	Ala	11e	His O	Gln	Thr	Ser	Gln 1099	Gly	Leu	Ala	Thr	Val 110		Lys	Ala	Leu
	Ala 110	Lys 5	Val	Gln	Asp	Val 1110	Val	Asn	Thr	Gln	Gly 1115	Gln S	Ala	Leu	Ser	His 1120
20	Leu	Thr	Val	Gln	Leu 1125	Gln 5	Asn	Asn	Phe	Gln 1130		Ile	Ser	Ser	Ser 1135	
	Ser	Asp	Ile	Tyr 1140	Asn)	Arg	Leu	Asp	Glu 1145	Leu 5	Ser	Ala	ązĄ	Ala 1150		Val
25	Asp	Arg	Leu 1155	Ile	Thr	Gly	Arg	Leu 1160		Ala	Leu	Asn	Ala 1165		Val	ser
	Gln	Thr 1170	Leu'	Thr	Arg	Gln	Ala 1175		Val	Arg	Ala	Ser 1180		Gln	Leu	Ala
30	Lys 1185	Asp	Lys	Val	Asn	Glu 1190	Cys	Val	Arg	Ser	Gln 1195		Gln	Arg	Phe	Gly 1200
	Phe	Cys	Gly	Asn	Gly 1205	Thr	His	Leu	Phe	Ser 1210		Ala	Asn	Ala	Ala 1215	
35	Asn	Gly	Met	lle 1220	Phē	Pħe	His	Thr	Val 1225		Leu	Pro	Thr	Ala 1230		G]u
	Thr	Val	Thr 1235	Ala	Trp	Ser		Ile 1240		Àla	Ser		Gly 1245		Arg	Thr
40	Phe	Gly 1250	Leu	Val	Val	Lys	Asp 1255	Val	Gln	Leu		Leu 1260		Arg	Asn	Leu
	Asp 1265	Asp	Lys	Phe	Tyr	Leu 1270	Thr	Pro	Arg		Met 1275		Gln	Pro	Arg	Val 1280
15	Ala	Thr	Ser	Ser	Asp 1285	Phe	Val	Gln		Glu 1290		Cys	Asp	Val	Leu 1295	

48

50

	Val	Asn	Ala	Thr 1300		Ile	Asp	Leu	Pro 1305		Ile	Ile	Pro	Asp 1310		Ile
5	Asp	Ile	Asn 1315		Thr	Val	Gln	Asp 1320		Leu	Glu	Asn	Phe 1325		Pro	Asn
	Trp	Thr 1330		Pro	Glu	Leu	Thr 1335		Asp	Ile	Phe	Asn 1340		Thr	Tyr	Leu
10	Asn 1345		Thr	Gly	Glu	Ile 1350		Asp	Leu	Glu	Phe 1355	Arg	Ser	Glu	Lys	Leu 1360
	His	Asn	Thr		Val 1365		Leu	Ala	Val	Leu 1370		Asp	Asn	Ile	Asn 1375	
15	Thr	Leu	Val	Asn 1380		Glu	Trp	Leu	Asn 1385		Ile	Glu	Thr	Tyr 1390		Lys
	Trp	Pro	Trp 1395		Val	Trp	Leu	Leu 1400		Gly	Leu	Val	Val 1405		Phe	Cys
20	Ile	Pro 1410		Leu	Leu	Phe	Cys 1415		Cys	Ser	Thr	Gly 1420	Cys)	Cys	Gly	Cys
	Ile 1425		Cys	Leu	Gly	Ser 1430		Cys	His	Ser	Met 1435	Cys	Ser	Arg	Arg	Gln 1440
25	Phe	Glu	Ser	Tyr	Glu 1445		Thr	Glu	Lys	Val 1450		Val	His			

Claims

- 30
- 1. A nucleic acid sequence encoding a polypeptide having one or more immunogenic determinants of a CCV spike protein.
- 2. A nucleic acid sequence according to claim 1, characterized in that the spike protein has an amino acid sequence shown in SEQ ID NO: 2, 4 or 6 or is a functional variant thereof.
 - 3. A nucleic acid sequence according to claim 2, characterized in that the nucleic acid sequence contains at least part of the DNA sequence shown in SEQ ID NO: 1, 3 or 5.
- 40 4. A recombinant vector molecule comprising a nucleic acid sequence according to claims 1-3.
 - 5. A recombinant vector molecule according to claim 4, characterized in that the nucleic acid sequence is operably linked to expression control sequences.
- 45 6. A recombinant vector virus harbouring the heterologous nucleic acid sequence according to claims 1-3.
 - 7. A host cell transformed with a nucleic acid sequence according to claims 1-3 or with a recombinant vector molecule according to claim 4 or 5, or infected with a recombinant vector virus according to claim 6.
 - 8. A process for the preparation of a polypeptide having one or more immunogenic determinants of a CCV spike protein which process comprises:
 - (a) culturing host cells according to claim 7 under conditions in which the nucleic acid sequence is expressed, and
 - (b) isolating the polypeptide from the culture.
 - 9. A vaccine for the protection of dogs against CCV infection or disease, characterized in that it comprises a recombinant vector virus according to claim 6, a host cell according to claim 7, or a polypeptide

50

prepared by the process according to claim 8, together with an acceptable carrier.

- 10. A process for the preparation of a CCV vaccine comprising the steps of culturing an infected host cell according to claim 7, collecting recombinant vector virus material, and formulating the material to a pharmaceutical preparation with immunizing activity.
- 11. A process for the preparation of a CCV vaccine comprising formulating a polypeptide prepared to the process of claim 8 according to a pharmaceutical preparation with immunizing activity.
- 10 12. A process for the protection of dogs against CCV infection comprising administering a vaccine according to claim 9 to a dog.

50

15

20

25

30

35

40

45

50

Figure 1




Figure 2a




Figure 2b





Figure 3





EUROPEAN SEARCH REPORT

EP 92 20 1136

	Citation of dominant with in	dication, where appropriate,	Relevant	CLASSIFICATION OF THE
stegory	of relevant pas		to claim	APPLICATION (Int. Cl.5)
	EP-A-0 295 057 (NORDEN	LABORATORIES, INC.)	1-12	C12N15/50
1	* Whole document, in par	rticular page 3, lines]	A61K39/215
ŀ	50-51, page 5 lines 51-	56 *]	G01N33/569
	50-51, page 5 (mas er			C07K15/00
۲	EP-A-0 376 744 (CALIFOR	NIA BIOTECHNOLOGY, INC)	1-12	
	* Whole document *			
.		-		
^	EP-A-0 138 242 (DUPHAR :	INTERNATIONAL RESEARCH	1,4,7,	
	B. V.)		9-10	
	* Whole document *			
		-		
Y	JOURNAL OF CLINICAL MIC	ROBIOLOGY	1-5,7-12	
	vol. 29, no. I, January	1991, US		
	I BAE ET AL.: 'Differs	entiation of transmissible	e	
	nastroenteritis virus f	rom porcine respiratory	1	
	coronavirus and other a	ntigenically related	İ	
	coronaviruses by using	cDNA probes specific for		
	the 5' region of the S	alycoprotein gene'	, ,	
	* Whole article *		1	
		_		
٧	EP-A-0 278 541 (DUPHAR	INTERNATIONAL RESEARCH	1-5,7-12	TECHNICAL FIELDS SEARCHED (Int. CL5)
	B. V.)			SEARCHED (III. C.S.)
	* Whole document *		İ	
		-		C12N
A	EP-A-0 344 872 (AMERICA	N HOME PRODUCTS	1	A61K
	CORPORATION)			C07K
	" Whole document *		i	
		-	1,4,6-12	
P,X,	WO-A-9 111 525 (THE UNI	VERSITY COURT OF THE	2,4,0-12	
D	UNIVERSITY OF GLASCOW)			ľ
	* Whole document *			
				
			1	
]	
			1	ļ
				1
	The present search report has b	een drawn up for all cinims		
	Place of searth	Date of completion of the sam		Examinat
	BERLIN	07 JULY 1992	JUL	IA P,
				- h-venting
_	CATEGORY OF CITED DOCUME	NTS T: theory or p E: earlier and	principle underlying the ent document, but pub) ished on, or
X : p2	rticularly relevant if taken alone	after the	iling cate	
₩	rticularly relevant if combined with an	other D: focument	cited in the application cited for other reasons	•
1 : br	and and the came and annual	L: eocument	CHAIN IN AMEDICAL	
≹o A∶te	cument of the same category chnological background m-written disclosure	I. : Wochillen	f the same patent fami	

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
COLOR OR BLACK AND WHITE PHOTOGRAPHS
GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
□ OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.